

Journal für

# Reproduktionsmedizin und Endokrinologie

– Journal of Reproductive Medicine and Endocrinology –

Andrologie • Embryologie & Biologie • Endokrinologie • Ethik & Recht • Genetik  
Gynäkologie • Kontrazeption • Psychosomatik • Reproduktionsmedizin • Urologie



## Heparin and Related Substances: Physiology and Pathophysiology in General and in Human Reproduction -

**Part 1**

Würfel W

*J. Reproduktionsmed. Endokrinol* 2016; 13 (2), 38-48

[www.kup.at/repromedizin](http://www.kup.at/repromedizin)

Online-Datenbank mit Autoren- und Stichwortsuche

Offizielles Organ: AGRBM, BRZ, DVR, DGA, DGGEF, DGRM, D-I-R, EFA, OEGRM, SRBM/DGE

Indexed in EMBASE/Excerpta Medica/Scopus

Krause & Pachernegg GmbH, Verlag für Medizin und Wirtschaft, A-3003 Gablitz



ENDO FERTI FORUM

ENDOKRINOLOGIE & FERTILITÄT  
FÜR KLINIK & PRAXIS

**20.-21. März 2026**

Universitätsmedizin Mainz

## Einladung zu unserer wissenschaftlichen Veranstaltung Endo-Ferti-Forum

Brücke(n) zwischen Unikliniken und Praxen an Rhein und Main(z)

– die aus dem bisherigen Format „Ferti Forum“ ab 2026 hervorgeht –



Freuen Sie sich auf spannende Vorträge und den lebendigen Austausch mit Kolleg:innen und Expert:innen aus Klinik und Praxis. Freitagabend laden wir Sie herzlich zu einem entspannten Empfang ein – eine perfekte Gelegenheit, Kontakte zu knüpfen und den Tag genussvoll ausklingen zu lassen.

Wissenschaftliche Leitung: Univ.-Professorin Annette Hasenburg, Dr. Susanne Theis, Universitätsmedizin Mainz, Sanitätsrat Dr. Werner Harlfinger, BVF Rheinland-Pfalz Dr. Rüdiger Gaase, BVF Hessen Dr. Klaus J. Doubek

Schirmherrschaften: Prof. Nicole Sänger, Uniklinik Bonn, Prof. Jan-Steffen Krüssel, Uniklinik Düsseldorf, Dr. Annette Bachmann, Uniklinik Frankfurt am Main, Prof. Christine Skala, Uniklinik Köln

Weitere Informationen  
& Anmeldung unter



# Heparin and Related Substances: Physiology and Pathophysiology in General and in Human Reproduction – Part 1\*

W. Würfel

Although heparin is usually associated with blood clotting and thrombosis prevention, its anti-coagulatory properties play an extremely significant role in clinical routine. However, it appears to have little importance in rheological homeostasis, particularly since heparin and related substances can also be found in simpler organisms without clotting mechanisms. In fact, heparin and heparan sulfate glycans are proteoglycans that have a long history in the development of species. Accordingly, their primary effects are electrostatic, not receptor-mediated, and they exhibit binding via their pronounced electronegativity. They exist in free form and as membrane-bound molecules. In physiological terms, heparin and heparinoids are immunological co-factors involved in inflammation and wound healing; they play significant roles in embryo development and organogenesis, serve as “homing factors,” and are involved at various points in carcinogenesis. They show diverse interactions with cytokines, growth factors and other mediators that are important in the implantation process and its control and continue to be so throughout pregnancy. It can be assumed that reproduction is a key focus of physiological functions; this would explain why heparin is so effective above and beyond its anti-coagulatory effects, e.g. in treating anti-phospholipid syndrome. To date no clear confirmation of the existence of a heparin deficiency syndrome has been possible, although initial results from studies concerning the general administration of heparin during pregnancy point in this direction. **J Reproduktionsmed Endokrinol Online 2016; 13 (2): 38–48.**

**Key words:** heparin, heparinoids, heparan sulfate glycans, history, measurement, human reproduction, rheology, endometrium, embryonic and fetal development, immunology, antiphospholipid syndrome, cancer

## ■ Introduction

Heparin is used routinely as a “blood thinner,” and heparin’s anti-coagulatory properties are doubtless its most prominent characteristic [1]. However, heparin and heparan sulfate glycans can also be found – in some cases, in high concentrations – in organisms without a blood clotting system [1], such as fresh-water clams [2], prawns [3], and other invertebrates [1]. This indicates on the one hand that the anti-coagulatory properties of heparin (or related substances) reflect only part of their physiological significance [4, 5], while also proving that this family of substances has an extremely long phylogenetic history [4, 5].

## ■ History of the Discovery of Heparin

Jay McLean (1890–1957), a 4<sup>th</sup>-semester student at John Hopkins University under Professor William Howell, had been assigned the task of extracting a stable thrombophilic substance based on cephalin prepared by Howell from rabbit brains [6, 7]. He used procedures already described by Erlandsen in 1907 for iso-

lating cuorin (from ox heart) [8], and by Baskoff for identifying heparophosphatides (from horse liver) [9]. However, the ether-based extraction performed by McLean in 1916 resulted in the isolation not of a thromboplastic, but an anti-coagulant substance, initially believed by McLean to be anti-thrombin [10]. Admittedly, in his memoirs he claims to have soon realized the substance was a heparophosphatide and cites this as the reason for its later name of heparin [6]. The name was first used in 1918, by which time McLean had left the university for financial reasons [11]; his work was continued under William Howell, primarily by Emmett Holt jr. [12]. However, the fat-soluble substance extracted from canine liver by Holt and named “heparin” naturally had a different biochemical makeup from McLean’s heparophosphatide. In 1922 the two scientists and their working group described a water-soluble polysaccharide [11], presenting a more advanced version in 1925 [13]. Clinical tests were performed at institutions including the Mayo Clinic and revealed frequent side-effects such as headache, fever and nausea [14]. Yet heparin was commercially available

from 1924, albeit without any significant medical role. In Canada, a group of scientists which formed around Charles Best from 1929 [15] published an improved procedure for isolating heparin from horse liver in 1933 (Arthur Charles and David Scott, 1933 [16–18]). This heparin, produced by Connaught Laboratories in Toronto, had significantly fewer side-effects, although still proving relatively toxic in individual cases [15]. Not until 1937 was a crystalline heparin produced, virtually free from side-effects and linked with the name of Gordon Murray [19, 20]. Its molecular structure was identified in 1935 by Eric Jorpes from the Karolinska Institute in Sweden, who had visited the Connaught Laboratories around 1929 [21]. One year later, the company Vitrum AB launched the first heparin for IV use, which was still relatively toxic [22]. The development of heparin preparations for human use that were largely free from side-effects took place in parallel in Canada and Sweden.

Jay McLean later also resumed his study of heparin, focusing on its clinical use in cases such as venous occlusion and endocarditis [23, 24].

\* Translated, updated and extended version from: Würfel W. Heparin und Heparinoide – was wissen wir über diese Substanzen? 1. Biochemie und gegenwärtige klinische Bedeutung. *gynäkol prax* 2013; 37: 205–15. With kind permission by Hans Marseille Verlag GmbH, Büro Wien.

Received: September 8, 2015; accepted after revision: February 2, 2016 (responsible Editor: Prof. Dr. W. Urdl, Graz)

From the Kinderwunsch Centrum München (KCM)

**Correspondence:** Prof. Dr. Dr. med. habil. Wolfgang Würfel, Kinderwunsch Centrum München (KCM), D-81241 München, Lortzingstraße 26; e-mail: info@ivf-muenchen.de

## ■ Molecular Structure

Heparin and its related substances such as heparan sulfate are proteoglycans, or “protein-carbohydrate hormones” (proteo-polysaccharides). These polysaccharides form linear chains in a polydisperse structure; the main constituents are pyranosyl-uronic acid (L-iduronic acid + D-glucuronic acid) + 2-amino glucosamine (Fig. 1), with variable acetyl or sulfo groups [25].

Heparan sulfates show far higher variability than heparin itself, with dominating glucuronic acid and reduced sulfo groups [26].

For this reason the level of sulfation is a popular method for distinguishing between heparin and heparan sulfates; however, no precise biochemical distinction has ever been successfully made.


Heparan sulfate proteoglycans (HSPG) are classified into three families based on their biochemical and/or physiological characteristics:

- Membrane-spanning HSPGs (e. g. syndecan [27], betaglycan [28] and CD 44 [29])
- Glycophosphatidylinositol-anchored HSPGs (e.g. glypicans [30])
- ECM (extracellular matrix) proteoglycans (e.g. agrin [31], perlecan [32] and collagen XVII [33]), which may be soluble or transmembrane (such as collagen XVII)

See also Figure 2.

Heparin and heparan sulfate glycans both belong to the family of glycosaminoglycans (GAGs) and have the highest electronegative charge known of any physiological molecules in our organism [4]: heparin has approx. 2.7 negatively charged groups per disaccharide (sulfo and carboxyl groups) whereas HSPGs have fewer than two. Accordingly, ion binding and not receptor binding is the focus of interaction with other molecules, and thus also of mediation of the effect. This makes clear that heparin/HSPG activity is strongly dependent on molecule size.

For example, a minimum of 17–18 saccharide groups are required for sufficient bioactivity response to thrombin (coagulant factor II). This does not apply to, say, ultra-low-molecular-weight (ulmw) heparins, so that their “general” bioactiv-



**Figure 1:** Monosaccharide components of heparin and heparinoids. The bracketed groups are relatively rare. SO<sub>3</sub> and NH<sub>2</sub> groups may be found as substituents in the D-glucosamines.

ity is lower than that of native heparin. However, the anticoagulation effect of ulmw heparins is only selective and this is often delivered via pentasaccharide, the “core ligand” (see below).

The high electronegative charge results in high binding affinity which extends to all proteins, particularly:

- Growth factors [35, 36]
- Chemokines and cytokines [37, 38]
- Lipoproteins [39]
- Receptor (protein) complexes [40]
- Proteases/esterases and their inhibitors [41, 42]
- Enzymes [43, 44]
- Viral coat proteins [45, 46]

An overview is given in Table 1.

As already mentioned, a type of ligand/receptor-mediated interaction exists, based on that of L-iduronic acid and its steric plasticity, creating a definable pentagonal structure: GlcNAc/NS(6S) – GlcA-GlcNS(3S,6S)-IdoA(2S)-GlcNS(6S) [48]. It is named DEFGH (the letters are historical abbreviations of the active groups or molecular positions in this pentasaccharide) [49], and is also known as high-affinity heparin (HAH). This pentasaccharide is essential for clinically significant interaction with activated coagulation factor Xa.

Where highly selective Xa suppression is desired, this pentasaccharide is actually

all that is needed – the rationale behind the develop of the synthetic “heparin” Fondaparinux (Arixtra®) [50].


Only around 30% of naturally occurring heparins function using HAH, while around 70% use ion binding [50].

## ■ Synthesis of Heparin and Heparan Sulfates

Heparin is present in humans in the liver (approx. 25 IU/100 mg), lungs, and muscular tissue, and at lower levels also in the intestine (intestinal mucous membrane), spleen and thymus gland [25]. Similar conclusions were reached by Charles and Scott in 1933 [17], who also found that heparin is virtually undetectable in serum – contrary to Howell’s assumptions [12]. Synthesis can be clearly detected in neutrophilous granulocytes and – especially – in mastocytes, i.e. immune system cells, and in endothelial cells [25, 51].

Synthesis of heparan sulfates can be proven for an extremely large number of cell types, which is only logical given their importance for membranes and the extracellular matrix (ECM) [51].

Intracellular synthesis occurs in the Golgi apparatus. The initial step is synthesis of a tetrasaccharide comprising glucuronic acid, 2 molecules of galactose and a molecule of xylose bonded to a core protein [52]. When chain formation is



**Figure 2:** The diagram shows transmembrane heparan sulfates each bound to protein complexes. Collagen XVII, which also has a transmembrane form, is not shown. Mod. from [4].

initiated, further saccharides are gradually added to form a polymer. In a third stage, the chain is modified, e.g. by sulfation or epimerization (into iduronic acid) [52].

The synthesis paths of heparin and heparan sulfates are thus virtually identical, making it harder to provide evidence of types of tissue where heparin synthesis does and does not take place. One theory, certainly worthy of serious consideration, states that synthesis takes place only in immunocompetent cells and in the endothelium (see below).

Unfractionated heparin (uf heparin) has a molecular weight of approx. 12–40 kilodaltons (kDa), with an average of approx. 14 kDa [50]. Uf heparins for clinical use are generally produced from bovine lung or porcine (intestinal) mucous membrane; biological activity of both organisms is similar, although the mean molecular weight of bovine heparins is slightly higher [53].

Low-molecular-weight heparin (lmw heparin), i.e. “fractionated” heparin, is produced from uf heparin (preferably porcine) by means of chemical or enzymatic depolymerization (heparinases, see below) and various fractionation methods [54]. Uf heparin already contains around 20–30% lmw heparins [53].

Low-molecular-weight heparins have better bioavailability and longer-lasting effects, and also cause fewer undesirable side-effects – presumably because they undergo far less (unspecific) binding with proteins, tissues and, primarily, blood cells (principally HIT-2, see below) [55]. Anticoagulation activity generates 80–120 anti-Xa units (i.e. U/mg or 35–45 anti-IIa U/mg [55]).

It is the uf heparins that show varying pharmacological and anti-coagulatory properties depending on their source material and production method. Anti-coagulatory units (IU) were defined to guarantee comparability of anti-thrombotic effects and express the in-vitro effect (5th International Standard for Unfractionated Heparin, NIBSC-Code 97/578). According to the standard, parenterally applied heparin must show specific anti-coagulatory activity of 150 IU/mg.

By contrast, anti-Xa activity is used as the basis for measuring effectiveness of lmw heparins (aXa-IU/mg); anti-IIa activity (aIIa-IU/mg) and the ratio of aXa/aIIa must also be stated [56].

A definition of biological activity is difficult, even from a historical perspective. The “Howell unit” was the first attempt, defining the quantity of uf heparin necessary to prevent coagulation of 1 ml of

feline blood for 24 h at 0 degrees, i.e. approx. 0.002 mg [57]. The United States Pharmacopoeia Unit (USP Unit) defines it differently, as the quantity of heparin necessary to cause only 50% of coagulation in 1 ml of ovine plasma in 1 h at 37 degrees Celsius [57].

## ■ Biodegradation of Heparin and Heparan sulfates

Heparan sulfates are broken down in the lysosomes. At present the assumption is a single heparanase (Hpa-1); the coded gene is located at chromosome 4 (q22) (approx. 40–50 kDa) [58]. A number of splicing variations exist [59]. Evidence of the presence of Hpa-1 is practically confined to lymphoid tissue, so that its effects can be assumed to be largely extracellular, e.g. in inflammation processes or during embryonal implantation [60]. Heparanase activity is also shown by CTAP-III (connective tissue activating peptide), present in thrombocytes [61], and by Hpa-2, shown to be present in non-lymphoid tissue [62]. The extent to which these are independent biodegradation pathways is the subject of scientific debate.

Numerous other enzymes are involved in biodegradation, including sulfatases, an iduronidase and a glucuronidase [60].

Hpa-1 is overexpressed in some human malignomas [63], with the degree of overexpression correlating to the negativity of the prognosis [64]. One reason could be that Hpa-1 biodegrades the heparan sulfates in basal membranes and thus promotes invasion of the malignoma [65].

A distinction must be made between heparanases and heparinases [66]; the latter were first identified in *Pedobacter heparinus* (previously *Flavobacterium heparinum*) and other sources [67]. 3 different heparinases – I, II and III (synonyms: heparin lyases or heparitinases). They are able to break down polymers at different points [66, 67].

In humans, heparin is primarily broken down in the liver (lysosomes), but also in the kidney, in a process described as uro-heparinase [67]. 20–30 % of heparin is excreted unaltered; the degradation products of heparin and heparan sulfates are excreted in the urine.

## ■ Coagulation

The anti-coagulatory effect of heparin is primarily based on activation of antithrombin III (AT III) [68], part of the family of serine protease inhibitors (serpins). They include

- heparin cofactor II (HC II),
- plasminogen activator inhibitor 1 (PAI-1),
- protease nexin-1 (PN1) and
- protein-C inhibitor (PCI, also known as PAI-3).

These serpins can bind to heparin or heparin-like glycosamin glycanes (GAG) [69] and this influence coagulation, although activation of AT III is the dominant factor in the effect of heparin.

Antithrombin III is an inhibitor of activated factor X (Xa) and activated thrombin (IIa) [69]. In addition, a significantly lower effect is exercised via HC II, which, however, only inhibits thrombin [70]. Although antithrombin III and HC II have very similar structures [71], AT III only binds to heparin, while HC II also binds to further heparinoid, i.e. GAGs, such as dermatan sulfate [72].

Antithrombin III inhibits further serine proteases including factor IXa, as well as factors XIa and XIIa and kallikrein. AT III has no effect on factor V and factor VIII [69].

Electropositive domains known as exosites mediate the electrostatic bond between heparin and antithrombin III [73]. They show concentrations of the aminoacids arginine and lysine [74]. The conformation changes upon binding with heparin, forming a complex that has higher potential for coagulation factor inhibition than antithrombin alone [71]. Relatively long-chain heparins are necessary to generate a sufficiently inhibitory effect, which also applies to HC II [75]. However, serpin activation by the pentasaccharide (HAH) is sufficient for the inhibition of factor Xa alone [76], forming the basis for the consideration of fondaparinux as already described (see above).

TFPI (tissue factor pathway inhibitor) drives a different mechanism of action of heparin [77]. This inhibitor is primarily synthesized in the vascular endothelium [78] and inhibits factors Xa and VIIa in a

**Table 1:** Summary of the main interactions of heparin and HPSG. Mod. from [47].

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |                                                                                                                                                                                                            |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Growth factors</b> <ul style="list-style-type: none"> <li>– FGF (fibroblast growth factor)</li> <li>– HB-EGF (heparin-binding epidermal growth factor)</li> <li>– EGF (epidermal growth factor)</li> <li>– IGF-2 (insulin-like growth factor)</li> <li>– TGF-<math>\beta</math> (transforming growth factor)</li> <li>– Hepatocyte growth factor</li> <li>– PGF-4 (platelet growth factor)</li> <li>– VEGF (vaso-endothelial growth factor)</li> <li>– G(M)-CSF (granulocyte (macrophage) colony stimulating factor)</li> </ul> | <b>Chemokines, further cytokines</b> <ul style="list-style-type: none"> <li>– IL-6, IL-8 (GP-130-cytokines)</li> <li>– IFN-<math>\gamma</math></li> <li>– SDF-1a (stromal-derived factor)</li> </ul>       |
| <b>Proteases/esterases and their inhibitors</b> <ul style="list-style-type: none"> <li>– Antithrombin (serpins), primarily III</li> <li>– HC II</li> <li>– Protein C inhibitor</li> <li>– PAI (plasminogen activator inhibitor)</li> <li>– Complement/complement inhibitors</li> <li>– Nexin-1</li> <li>– TFPI (tissue factor pathway inhibitor)</li> <li>– Secretory leukocyte protease inhibitor</li> </ul>                                                                                                                      | <b>Receptor (protein) complexes</b> <ul style="list-style-type: none"> <li>– EGF receptor</li> </ul>                                                                                                       |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <b>Enzymes</b> <ul style="list-style-type: none"> <li>– Factors Xa, IIa (thrombin) IXa</li> <li>– Neutrophil elastase</li> <li>– Kathepsin G</li> <li>– Superoxide dismutase</li> </ul>                    |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <b>Viral envelope proteins</b> <ul style="list-style-type: none"> <li>– HIV envelope protein GB-120</li> <li>– HSV envelope proteins B (gB), gC and gD</li> <li>– Dengue virus envelope protein</li> </ul> |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <b>Lipoproteins</b> <ul style="list-style-type: none"> <li>– Apolipoprotein E</li> <li>– Lipid-binding proteins</li> </ul>                                                                                 |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <b>Further proteins</b> <ul style="list-style-type: none"> <li>– Malaria proteins</li> <li>– Endotoxin</li> <li>– Annexin V</li> </ul>                                                                     |

two-stage reversible reaction [79]. Like other GaGs, heparin can release TFPI [80]. Around one-third of heparin's anti-coagulatory effect is determined by TFPI [81], causing higher TFPI levels to show in serum after heparin treatment [77, 81].

Heparin is known to trigger forms of thrombocytopenia, namely heparin-induced Type 1 and Type 2 thrombocytopenia (HIT-1 and HIT-2). HIT-2 can take on life-threatening forms by inducing thrombosis and embolism [82] (Fig. 2).

The pathophysiological focus here is the increased release of platelet factor 4 (PF-4) from thrombocytes and subsequent formation of a PF-4/heparin complex [82]. HAH plays a particularly significant role in platelet activation, as does the density of the receptor status on the thrombocytes [83]. A major role is played by heparin's high electronegativity, while PF-4 shows high electropositivity [84]. This electrostatic bond also explains why such complex formation is found more frequently in long-chain heparins (uf) than short-chain heparins (nm) or even pentasaccharide alone [82].

A decisive element in the further pathophysiological cascade is the formation of antibodies (IgG) against the PF-4/heparin complex, albeit virtually exclusively directed against PF-4 [85]. Once such antibodies form, they in turn activate the

thrombocytes. This increases the release of PF-4, which can antagonize heparin and bind HSPGs to cell membranes [84, 86].

Differential diagnosis of both HITs is clinically significant as the consequences for treatment differ (Tab. 2). HIT-2 occurs far more rarely in fractionated, i.e. low-molecular heparins [88, 89], and is extremely unusual during pregnancy; accordingly, there is only one publication on the subject [89].

Protamine is a mixture of strongly alkaline peptides and has a discrete anti-coagulatory effect. However, formation of a heparin complex is essential to trigger this effect. Upon heparin complex formation it inhibits heparin-induced anti-Xa effect completely, and anti-IIa effect partially. Fondaparinux remains unaffected [90].

Protamines are produced from the sperm or roe of some salmonids. The neutral water-soluble salts such as protamine sulfate and protamine hydrochloride are used [90]. The effect develops rapidly (5-15 min). Given the slight anti-coagulatory effect, overdosing may cause haemorrhage [91].

No essential physiological significance of heparin for coagulation has yet been proved [92, 93], despite its frequent use as an anti-coagulant. Heparins are cur-

**Table 2:** Heparin-induced thrombocytopenias (HIT). Mod. from [87, 88].

|                     | HIT-1                                          | HIT-2                                                                           |
|---------------------|------------------------------------------------|---------------------------------------------------------------------------------|
| <b>Cause</b>        | Direct thrombocyte activation by heparin       | Formation of antibodies (et al.), PF-4 activation                               |
| <b>Start</b>        | 1–2 days after 1 <sup>st</sup> dose of heparin | 5–14 days after 1st dose of heparin, or earlier if dose administered previously |
| <b>Thrombocytes</b> | Rare < 100,000/ml                              | Decline > 50% of starting value                                                 |
| <b>Frequency</b>    | Approx. 10%                                    | Approx. 1%, rarer in pregnancy                                                  |
| <b>Clinical</b>     | Generally asymptomatic                         | Possible thromboembolic vascular occlusions, embolism                           |
| <b>Treatment</b>    | None                                           | Discontinue immediately, switch to alternative anticoagulant where applicable   |

rently assumed to be non-essential for physiological hemostasis [94]. The same appears to apply to HSPGs, which are generally membrane-bound and assumed to have a local effect [95]. Given the diverse and, in some cases, widely differing (co-) effects, the term “multivalent biomodulators” is applied to heparin and GaGs [96].

## ■ Immune System

Significant levels of heparin synthesis takes place in mastocytes. Synthesized heparin is stored in granules in the mastocytes and, together with histamine, is their most important (exocrine) synthesized product [96, 97]. Synthesis has also been shown to occur in neutrophil granulocytes [92, 93].

Mastocytes are found in high concentrations near blood and lymph vessels, along nerve cells, and particularly in mucous membranes and skin [93].

An often-repeated theory is that heparin synthesis only exists in mastocytes and endothelial cells and that the heparin content of e.g. lung or liver tissue is derived from these cells [98]. This is, however, not the case, as mastocytes are virtually absent from mucous membranes in various intestinal sections where heparin is found in relatively high concentrations [100].

The complement system, comprising approximately 25 different proteins [100], is an important partner for the effect of heparin. The main task of the complement system is opsonization, i.e. interaction with the surface of pathogens to enable phagocytes to destroy them. In addition, the complement system can trigger general inflammatory reactions,

may develop chemokine effects triggered by individual fragments, and may also attack bacterial envelopes [100, 101].

The complement system can be activated in three ways [101]:

- the classic method,
- the lectin method,
- the alternative method.

The classic method focuses on activation of glycoproteins C1-C9 by IgG or IgM antibodies, while IgA, IgE and IgD antibodies are ineffective. Another possibility is activation by direct binding of C1q to pathogen surfaces or to collagen or CRP [102].

In the lectin method, mannose-binding lectin (MBL) binds to mannose or glycans on the pathogen surface and thus activates proteases MASP-1 to -3, triggering the same cascade reaction as the classic method [100, 102].

The alternative method, also known as the properdin method, is activated by degradation of relatively unstable factor C3 into C3a and C3b. C3a has a chemotactic effect as an inflammatory anaphylatoxin. With other factors such as factor B, Cb3 can form the C3 convertase of the alternative method and trigger cascade reactions in the presence of pathogenic surfaces [101].

Heparin leads to partial inhibition of the complement system and its activation in both the classic activation method [103] and the properdin method [104]. The focus is on C3, which is of central importance in all activation paths [104] even though other complement factors [105, 106] such as C2, C4 or Hf are the target of the inflammation.

In addition, binding activity of specific proteins involved, e.g. C4 binding protein (decreases) or vitronectin (increases) is modulated [107, 108].

Individual factors of the complement system are also known to be activated. However, the inhibitory effects predominate overall, so that the effect of heparin on the complement system can best be characterized as immune-suppressive, as was identified as early as 1929 [109].

Formation of free radicals or reactive oxygen species (ROS) is also inhibited [110, 111], as is synthesis of pro-inflammatory cytokines in leukocytes, primarily neutrophil leukocytes [112], as well as in mononuclear cells and macrophages. The most important effect is inhibition of TNFα synthesis and release; in addition, heparin is a potent antagonist of the endotoxin [113, 114]. A further noteworthy effect is the inhibition of migration in eosinophil granulocytes [115].

Heparin therefore has anti-inflammatory properties, as has been shown in clinical studies. Positive therapeutic effects are found for e.g. rheumatoid arthritis [116], allergic bronchial asthma, allergic rhinitis [117], ulcerating colitis and Crohn's disease [118].

Heparin's somewhat general anti-inflammatory properties are countered by specific virostatic, bacteriostatic and even bacteriocidal effects caused by heparin's powerful negative charge. Heparin therefore binds to many proteins, including pathogenic proteins such as HIV envelope protein [119], the envelope protein of the herpes simplex virus (HSV) [120] and the envelope protein of the Dengue virus [121]; in the latter, the HAH complex evidently plays an important part in binding. There is now good evidence for heparin's anti-malarial activity [122, 123].

In this context HSPGs must be evaluated differently; for example, syndecan-3, a membrane-bound HSPG (see below), has a supporting effect in binding and cell transmission of the HI virus (HIV) and filoviridae [124, 125]. However, this need not result in aggravation of the disease, as it is perfectly possible that improved intracellular absorption of the viruses actually helps to eliminate them more rapidly [126].

Mastocytes play a major role in the bacteriostatic and bacteriocidal effect of heparin. They are shown to occur in many organs in contact with the external environment – such as lungs and dermis, where they occur in proximity to blood vessels, nerves and hair follicles – often in high concentrations [overview in 127]. Mastocyte activation is effected by bacteria or bacterial excretory products via specific receptors, e.g. CD 85 or toll-like receptors (TLR) 2, 4, 6, 7, 9) or inflammation mediators such as complement factors, endothelin-1 and neurotensin.

Mastocytes release various cytokines (e.g. TNF $\alpha$ ), enzymes and proteases (e.g. chymase, tryptase or carboxypeptidase) and/or chemokines, but primarily histamine and heparin. Heparin in particular establishes communication with other immunocompetent cells and thus influences differentiation of dendritic cells from monocytes [128].

Mastocytes are therefore extremely important for combating bacterial and parasitical infections (pruritis) and are closely involved in the innate immune system (cf. TLRs, see above). Heparin and histamine promote circulation in inflamed areas by vasodilation and anti-coagulation (flushing).

Mastocytes are also able to limit inflammatory reaction once triggered. This is useful given that proliferating and prolonged inflammation has a destructive potential that is hazardous to the organism. Heparin also appears to be extremely significant here [cf. 127].

Heparin thus possesses a pronounced immunomodulating effect.

It develops intrinsic effects in warding off infections, and communicates with various cells involved in specific immunodefence, e.g. dendritic cells. Together with other substances such as histamine, heparin ensures good local perfusion in the initiation and maintenance of inflammation; however, given its general anti-inflammatory properties it also plays a role in containing inflammation.

## ■ Cell Proliferation and Cellular Differentiation

HSPGs (and, to a lesser extent, heparin) in particular interact with a variety of

growth factors and hence modulate their effect on cell growth, cell differentiation, and thus cell development [129]. The majority of studies in existence focus on FGF (fibroblast growth factor) and VEGF (vascular endothelial growth factor) [130].

In their inactive forms, both these growth factors can be identified on the surface of endothelial cells or in the extra-cellular matrix, where they are bound to HSPGs [131]. This bond is broken by physiological or pathological stimulus, whereupon the growth factors form complexes with the HSPGs; these complexes significantly increase the binding affinity of the growth factors to their receptor [132].

This was particularly clearly shown for FGF-2. To date 22 different FGF variants have been identified, which differ – in some cases, considerably – in terms of their activation and receptor affinity [133].

The HAH complex with a minimum of one iduronic acid molecule in the 2-O position is critically important for this binding [134]. However, longer-chain HS fragments are necessary to initiate contact with the receptor [135]. This explains why some growth factor-HS complexes have an activating effect and others are inhibitory [135]. The same applies to heparin binding and the biological activity it triggers [136], for which reason the effects of unfractionated heparins and fractionated or short-chain heparins differ (see above).

Similar mechanisms have been shown for further growth factors [overview in 137], e.g.

- members of the EGF (epidermal growth factor) family,
- PDGF (platelet-derived growth factor),
- TGF- $\beta$  (transforming growth factor) 1 and 2 and binding protein,
- HGF (hepatocellular growth factor),
- IGF-2 (insulin-like growth factor) and IGF binding proteins 3 and 5,
- G(M)-CSF (granulocyte-macrophage colony-stimulating factor) and
- heparanases.

They are essential for the breakdown of heparin and HSPGs (see above). However, heparin and some HS fragments also have an intrinsic effect on these enzymes

[138], whereby increasing concentrations of heparin result in inhibition of heparanase activity [139]. This means that heparin bioactivity does not increase in a linear fashion, but exponentially from a specific concentration.

Direct influence on cell growth thus exists. Growth inhibition of smooth muscle cells has been shown both in culture medium [140] and directly on the endothelial wall [141, 142].

The fact that heparin binds specifically to smooth muscle cells has given rise to the question of whether a specific heparin receptor exists [142, 143]. It is assumed that no such receptor exists and that in fact, heparin and various HSPG fragments are capable of activating intracellular signal transduction in a similar way to extra-cellular activation (of growth factors, proteases, cytokines etc.).

Proven interactions include the following:

- Suppression of serum and glucocorticoid-regulated kinase (SGK), an early gene response to cell proliferation [144],
- Selective regulation of tyrosine kinase, a key protein in e.g. angiogenesis [145],
- Inhibition of mitogen-activated protein kinases [146],
- Inhibition of calcium/calmodulin-regulated protein kinase II [147].

Here too, the structure of the heparin or HSPGs is critically important; many biological effects can only be shown for longer-chain (generally unfractionated) heparins and heparinoids, not for their short-chain counterparts [148]. In turn, contrasting effects can accordingly be observed, i.e. activation instead of inhibition and vice versa [149].

This modulating influence on cell growth and cell differentiation shows that heparin plays an important role here. This is proven for e.g. cell repair processes with cicatricial build-up [150], embryo implantation followed by pregnancy [4, 151] and malignant infiltration [152, 153].

The important role of numerous membrane-bound HSPGs, such as syndecane [154], glypican [155] and perlecan [156], has also been shown for these processes.

## ■ Membranous Heparan Sulfate Glycans

Membranous HPSGs have a different physiological significance than soluble HPSGs.

In general, it may be assumed that membrane-bound structures target cell interaction and regulation of substrate metabolism between extra-cellular and intra-cellular space and vice versa. In addition, it can be assumed that membrane-bound molecules extending far into extracellular space have an affinity for various extra-cellular ligands.

The following groups or molecules are distinctly defined:

- Syndecans
- Glypicans
- Betaglycans
- CD44

### Syndecans

Syndecans are transmembranous HPSGs anchored with a CoR protein and associated with chondroitin sulfate chains (see Fig. 2). Four members of this HSPG family are currently known (1–4), specified in their sequence of cloning [157]. While the transmembranous and intra-cellular part is extremely similar throughout all species, the extracellular part varies considerably [158].

Distribution in tissue varies, in some cases widely:

- Syndecan-1 is primarily found in epithelial tissues, but also in mesodermal tissue e.g. skeletal muscle [157, 159].
- Syndecan-2 primarily occurs in mesenchymal cells/tissues, e.g. fibroblasts and hepatocytes [160].
- Syndecan-3 is primarily found in nerve and muscle tissue [161].
- Syndecan-4 is found the most frequently in the organism and occurs in epithelial and endothelial cells, e.g. fibroblasts, myocytes, chondrocytes and nerve cells [157, 162].

Syndecans bind to various substances, particularly growth factors (see above) and primarily FGF-2, but also TGF- $\beta$  (syndecan-2) [163], EGF, VEGF, and HBF [157]. Their actual function is that of a co-receptor (see: FGF binding), principally for g-protein coupled receptors; a further function is cell adhesion

(primarily syndecan-1), with noteworthy interactions with integrin and fibronectin [164].

Syndecans are also important during phylogenesis [165]. Their presence can first be shown from the 16-32-cell stage of the preimplantation embryo, when they are found on all cells of the inner cell mass (ICM) of the blastocyst and they are among those responsible for cell adherence [166]. They are also important in advanced embryo development, particularly in development of epithelial tissue.

The same applies to neurogenesis, primarily with respect to syndecan-2 and syndecan-3 [167], i.e. development of neurons (from neural stem cells) and neural synapses [168]. Syndecans also play a role in muscular development, particularly syndecan-4, which partly develops its effect through growth factors FGF-2 and TGF- $\beta$  [169]. Myogenic satellite cells, found between the basal membrane and the sarcolemma, are crucial in this context. They are activated in proliferation processes including repair. This results in expression of syndecan-3 and -4 [170] at all stages of life, including embryos [171].

### Glypicans

Glypicans are not transmembranous HPSGs, but are anchored to the cell surface by the GPI (glycosylphosphoinositol) anchor [172].

Six different glypicans (1–6) have been identified in vertebrates, with widely varying expression in tissues:

- Glypican-1 is primarily found in the brain, kidneys and skeletal musculature [173].
- Glypican-2 is important for the development of the nervous system [174].
- Glypican-3 is primarily found in mesodermal tissue and is strongly expressed during intestinal development [175].
- Glypican-4 primarily occurs in blood vessels [176].
- Glypican-5 occurs in brain and kidney tissue [177].
- Glypican-6 is the most frequently found, particularly in adult ovarian tissue, in fetal and adult kidneys, in smooth muscle tissue, in intestinal mesenchymal cells, in the lungs and in the teeth [178].

Similarly to syndecans, glypicans play an important role in regulating cell growth, e.g. in inducing cell apoptosis [179].

Although many aspects of the different interactions are still unclear, regulation of IGF-2 by glypicans, primarily glypican-3, appears to be important [173, 175, 178]. This is demonstrated by e.g. mutations in glypican-3 that can lead to excessive cell growth [180]. Here Simpson-Golabi-Behmel syndrome (SGBS) should be mentioned, characterized by features including pre- and post-natal overgrowth [181]. In this respect it is very similar to Beckwith-Wiedemann syndrome, and indeed both syndromes involve over-expression of IGF-2. This forms the basis for the theory that this syndrome is also caused by glypican-3 insufficiency (resulting from polymorphism) [180].

Syndecan-1 and glypican-1 have a synergistic effect on musculature by means of activation of FGF-2, resulting in increased cell proliferation with simultaneous inhibition of differentiation [173].

Like many other growth factors, FGF-2 is also involved in neurogenesis, primarily where stem cell differentiation is concerned [182]. Glypicans are also involved here; the precise nature of the interactions is unclear [182]. High expression of glypican-4 in the ventricular zone of the developing brain has been proven, particularly in neural cells with stem cell properties. After completion of differentiation, however, this expression recedes to the point of negligibility [183], although other glypicans (e.g. -1, -2, -5) show high expression in differentiated and post-mitotic neurons [174, 177].

### Beta Glycan

Unlike alpha glycans, beta glycan (see Fig. 2) is an HSPG which has transmembrane localization, a CoR protein and an extra-cellular domain.

Beta glycan is basically a co-receptor for members of the TGF- $\beta$  superfamily [184], particularly TGF- $\beta$ 2. Its effect does not result from its own intrinsic activity but from increased receptor affinity for TGF- $\beta$  at both its receptors (T $\beta$ RI and T $\beta$ RII) [185]. For this reason, beta glycan is also known as T $\beta$ -receptor-III (T $\beta$ RIII) [186].

The CoR protein shows particularly high affinity for members of the TGF- $\beta$  superfamily; HS side chains can also bind alkaline FGF (bFGF) [185].

Beta glycan also exists in soluble form in serum and the ECM, albeit in low concentrations. Unlike the membrane-bound beta glycans, this soluble type has an inhibitory effect on the TGF- $\beta$  family [187].

### CD44

CD44 is a glycoprotein occurring on cell surfaces. It exists in various forms that all originate from a single gene, [188] and are thus splicing variants (CD44v and the longer form of CD44E [on some epithelial cells]) [189]. The various (iso-) forms show differing concentrations of heparan, chondroitin and keratan sulfate [188, 189].

The physiological function of the CD44 family is primarily cell-to-cell adhesion. They can thus bind collagens and osteopontin, interact with metalloproteinases (MMP) and in some cases bind growth factors, albeit with low affinity [189, 190].

HCELL (e-selectin/l-selectin ligand of hematopoietic cells) is a sialated and glycosylated variant with functions including a “homing receptor” for bone marrow, i.e. (re-)directs hematopoietic and mesenchymal stem cells there [191]. It also appears to be important in carcinomas [192], particularly types of leukemia, in which it was first shown to occur on blast cells [193]. (Over-) expression on endometrial cells, particularly in foci of endometriosis, has also been shown. This higher expression is designed to promote peritoneal adherence [194].

### ■ Pericellular HSPGs

Unlike the HSPGs previously described, which are primarily limited to cell surfaces, pericellular HSPGs do not have CoR protein, are not anchored to cell membrane and are thus mainly found in the ECM, often in close proximity to cells or their membranes (“pericellular”).

They include perlecan, agrin and collagen XVIII.

### Perlecan

Perlecan takes its name from its appearance under an electron microscope, where it resembles a string of pearls.

Perlecan is a relatively widespread HPSG and has been identified in virtually all basal membranes [195]. It forms reticular structures with a profound filtering effect (strongly negative charge), e.g. in the kidneys.

Perlecan is coded by the HSPG2 gene. It comprises a core protein with approx. 660 KDa, has a high negative charge from residual sulfate, and has five domains which differ widely in some respects [196–198]:

- Domain I has HSPG chains – a domain for sperm protein enterokinase.
- Domain II contains receptors including one LDL receptor.
- Domain III has close similarity to the short arm of laminin A.
- Domain IV contains neural cell adhesion molecules and binds to nidogenes, fibronectin and heparin.
- Like domain IV, domain V also has molecular arrangements which correspond to EGF and bind to nidogenes, fibulin-2 and heparin.

Perlecan is crucial to normal embryonic growth [199]. Analyses of knock-out mice have shown that perlecan deficiency causes abnormal brain development and relatively premature mortality from skeletal dysplasia [200].

Development of the embryonic myocardium is a similar case [201]. Perlecan deficiency causes holes to form in the myocardium [201].

In both developing embryos and adults, perlecan deficiency prevents correct localization of acetylcholinesterase at neuromuscular junctions [202] yet without affecting actual acetylcholinesterase synthesis [203]. It can therefore be assumed that perlecan plays an important role in localizing and concentrating this vital neurotransmitter [204].

In combination with syndecan-1, perlecan anchors the lipoprotein lipase which splits triglycerides from chylomicrons. Given this mechanism, both HSPGs are involved in the lipid transport of chylomicrons to the epithelium [159, 199].

Mutations in the HSPG2 gene (1p36) are the cause of Schwartz-Jampel syndrome (chondrodystrophic myotonia) [205], dyssegmental dysplasia, Silverman-

Handmaker type [206] and probably also Roland-Debuquois syndrome [205, 206]. These overall extremely rare, but severe disorders differ in their level of perlecan synthesis, which is virtually absent in dyssegmental dysplasia but relatively high in Roland-Debuquois syndrome. The different perlecan synthesis rates would also correlate with the different clinical manifestations and their severity.

### Agrin

Agrin is a larger proteoglycan which, like syndecan and glypican, contains chondroitin and has three different binding sites for heparan sulfate [207]. It is coded by the AGRN gene [208].

Agrin derives its name from the synthesis or clustering of the acetylcholinesterase receptor (AChR) and aggregation of AChR during synaptogenesis, and from the importance of this proteoglycan in these processes [209, 210]. On the one hand, agrin inhibits cell overgrowth in the development of the nervous system, both peripheral and central (e.g. hippocampus, retina) [210]; on the other, agrin deficiency in knock-out mice impairs synapse development extensively or completely [211].

The same evidently applies to the MuSK receptor (receptor for muscle-specific kinase); agrin also binds to it and is thus responsible for (re-)phosphorylation [212]. Further muscle proteins to which agrin binds include laminin and dystroglycan.

### Collagen XVIII

Collagens (from the Greek for “glue-producing”) are structural proteins primarily found in the ECM. They are composed of  $\alpha$ -helical peptide chains (600–3000 amino acids) which are, unusually, left-handed and join to form a right-handed triple helix or superhelix (“coiled coil”).

29 types of collagen are currently known (I–XXIX), classified as fibrillar, reticulate, facit and short-chain, as well as collagens with anchoring fibrils and transmembrane collagens [213].

Collagen XVIII is a further HPSG for which 8 different glycosylation variations have been described.

Collagen XVIII is primarily found in endothelial and epithelial basal membrane

[214]. Individual glycosylation variations show anti-angiogenetic and anti-tumorous properties [215], primarily endostatin [215]. This is a split product of collagen XVIII with molecular weight of approx. 20 kDa, which is also endogenically effective as an angiogenesis inhibitor (198). Clinical studies have shown benefits from combining recombinant (rh) endostatin with chemotherapeutic agents in treatment of various types of malignoma [217, 218].

## Conflict of Interest

The authors have no conflict of interest to disclose.

## References

- Garg HG, Linhardt RJ, Hales CA (eds). Chemistry and biology of heparin and heparansulfate – preface. Elsevier, Oxford 2005; VII–VIII.
- Hovingh P, Linker A. Glycosaminoglycans in Anodonta californiensis, a freshwater mussel. Biol Bull 1993; 185: 263–76.
- Dietrich CP, Paiva JF, Castro RA, Chavante SF, Jeske W, et al. Structural features and anticoagulant activities of a novel natural low-molecular-weight heparin from the shrimp Penaeus brasiliensis. Biochem Biophys Acta 1999; 1428: 273–83.
- Nelson SM, Greer IA. The potential role of heparin in assisted conception. Hum Reprod Update 2008; 14: 623–45.
- Lever R, Page CP. Novel drug development opportunities for heparin. Nat Rev Drug Discov 2002; 1: 140–8.
- McLean J. The discovery of heparin. Circulation 1959; 19: 75–8.
- Howell WH. The factors concerned in the process of the clotting of blood. Therapeutic Gazette 1912; 36: 95–8.
- Erlandsen A. Untersuchungen über die lecitinhaltigen Substanzen des Myocardiums und der quergestreiften Muskeln. Hoppe-Seyler's Ztschr physiol Chemie 1907; 51: 71–155.
- Baskoff A. Über das Jecorin und andere lecitinhaltige Produkte der Pferdeleber. Hoppe-Seyler's Ztschr physiol Chemie 1908; 57: 395–460.
- Böttiger LE. The heparin story. Acta Med Scand 1987; 222: 195–200.
- Wardrop D, Keeling D. The story of the discovery of heparin and warfarin. Brit J Haematol 2008; 141: 757–63.
- Howell WH, Holt E jr. Two new factors in blood coagulation – heparin and pro-Thrombin. Am J Physiol 1918; 47: 328–41.
- Howell WH. The purification of heparin and its presence in blood. Am J Physiol 1925; 17: 553–62.
- Mason MC. A note on the use of heparin in blood transfusion. J Lab Clin Med 1924; 10: 203–6.
- Rutty CJ. Miracle blood lubricant: Connaught and the story of heparin, 1928–1937. Contact 1996; 9: 4–8.
- Charles AF, Scott DA. Studies on heparin I: the preparation of heparin. J Biol Chem 1933; 102: 425–9.
- Charles AF, Scott DA. Studies on heparin II: heparin in various tissues. J Biol Chem 1933; 102: 431–5.
- Charles AF, Scott DA. Studies on heparin III: the purification of heparin. J Biol Chem 1933; 102: 437–48.
- Murray GW, Jacques LB, Perett TS, Best CH. Heparin and the thrombosis of veins following injury. Surgery 1937; 2: 163–87.
- Murray GW. Heparin in thrombosis and embolism. Br J Surg 1940; 27: 324–36.
- Jorpes E. The chemistry of heparin. Biochem J 1935; 29: 1817–20.
- Jorpes EJ. Heparin: a mucopolysaccharide and an active antithrombotic drug. Circulation 1959; 19: 87–91.
- McLean J, Meyer BBM, Griffith JM. Heparin in subacute bacterial endocarditis. Reduction of cases and critical review of the literature. J Am Med Assoc 1941; 117: 1870–5.
- McLean J, Johnson AB. Gangrene following fracture treated with heparine, papaverine, and intermittent venous occlusion. Surgery 1946; 20: 324–36.
- Casu B, Lindhal U. Structure and biological interactions of heparin and heparansulfate. Adv Carbohydr Chem Biochem 2001; 57: 159–66.
- Linhardt RJ. Heparin: structure and activity. J Med Chem 2003; 46: 2551–64.
- Carey DJ. N-Syndecan: structure and function of a transmembrane heparin sulphate proteoglycan. Perspect Dev Neurobiol 1996; 3: 331–46.
- Andres JL, Stanley K, Cheifetz D, Massagué J. Membrane-anchored and soluble forms of betaglycan, a polymorphic proteoglycan that binds transforming growth factor-beta. J Cell Biol 1989; 109: 3137–45.
- Goodison S, Urquidí V, Tarin D. CD 44 cell adhesion molecules. Mol Pathol 1999; 52: 189–96.
- Filmus J. Glypican in growth control and cancer. Glycobiology 2001; 11: 19R–23R.
- Kallunki P, Eddy RL, Byers MG, Kestilä M, Shows TB. Cloning of human heparin sulphate proteoglycan core protein, assignment of the gene (HSPG2) to 1p36.1-p35 and identification of a BamHI restriction fragment length polymorphism. Genomics 1991; 11: 389–96.
- Tsen G, Halfter W, Kröger S, Cole GJ. Agrin is a heparin sulphate proteoglycan. J Biol Chem 1995; 270: 3392–9.
- Franzke C, Bruckner P, Bruckner-Tuderman L. Collagenous transmembrane proteins: recent insights into biology and pathology. J Biol Chem 2005; 280: 4005–8.
- Petitou M, Hérault JP, Bernat A, Driguez PA, Duchaussoy P, et al. Synthesis of thrombin inhibiting heparin mimetics without side effects. Nature 1999; 398: 417–22.
- Pillai SB, Turman MA, Besner GE. Heparin-binding EGF-like growth factor is cryoprotective for intestinal epithelial cells exposed to hypoxia. J Pediatr Surg 1998; 33: 973–9.
- Liang A, He X, Du Y, Wang K, Fung Y, Lin B. Quantitative investigation of the interaction between granulocyte-macrophage colony-stimulating factor and heparin by capillary zone electrophoresis. J Sep Sci 2006; 29: 1637–41.
- Gutiérrez G, Sarto A, Berod L, Canellada A, Gentile T, et al. Regulation of interleukin-6 fetal placental levels could be involved in the protective effect of low-molecular weight heparin treatment on murine spontaneous abortion. Am J Reprod Immunol 2004; 51: 160–5.
- Call Dr, Remick DC. Low molecular weight is associated with greater cytokine production in a stimulated whole blood model. Shock 1998; 10: 192–7.
- Gigli M, Consonni A, Ghiselli G, Rizzo V, Naggi A, Torri G. Heparin binding to human plasma low-density lipoproteins: dependence on heparin sulphate degree and chain length. Biochemistry 1992; 31: 5996–6003.
- Aviezer D, Yayon A. Heparin-dependent binding and autophosphorylation of epidermal growth factor (EGF) receptor by heparin-binding EGF-like growth factor but not by EGF. Proc Natl Acad Sci USA 1994; 91: 12173–7.
- Makrides SC. Therapeutic inhibition of the complement system. Pharmacol Rev 1998; 50: 59–88.
- Gettins PG. Serpin structure, mechanism, and function. Chem rev 2002; 12: 4751–804.
- Ermoloff J, Duranton J, Petitou M, Bieth JG. Heparin accelerates the inhibition of cathepsin G by mucus proteinase inhibitor: potent effect of O-butyrylated heparin. Biochem J 1998; 15: 1369–74.
- Brown RA, Lever R, Jones NA, Page CP. Effect of heparin and related molecules upon neutrophil aggregation and elastase release in vitro. Br J Pharmacol 2003; 139: 845–53.
- Lin YL, Lei HY, Lin YS, Yeh TM, Chen SH, Liu HS. Heparin inhibits dengue-2 virus infection of 5 human liver cell lines. Antiviral Res 2002; 56: 93–6.
- de Witte L, Bobardt M, Chatterji U, Degeest G, David G, Geijtenbeek TBH, Gallay P. Syndecan-3 is a dendritic cell-specific attachment receptor for HIV-1. Proc Natl Acad Sci 1007; 104: 19464–69.
- Munoz EM, Linhardt RJ. Heparin-binding domains in vascular biology. Arterioscl Thromb Vasc Biol 2004; 24: 1549–57.
- Lindhal U, Bäckström G, Thunberg L, Leder IG. Evidence for a 3-O-sulfated D-glucosamine residue in the antithrombin-binding sequence of heparin. Proc Natl Acad Sci USA 1980; 77: 6551–5.
- Atha DH, Stephens AW, Rimon A, Rosenberg RD. Sequence variation in heparin octasaccharides with high affinity from antithrombin III: Biochemistry 1984; 23: 5801–12.
- Rabenstein DL. Heparin and heparin sulphate: structure and function. Nat Prod Rep 2002; 19: 312–31.
- Esko JD, Selleck SB. Order out of chaos: assembly of ligand binding sites in heparan sulfate Annu Rev Biochem 2002; 71: 435–71.
- Perrimon N, Bernfield M. Specificities of heparan sulphate proteoglycans in developmental processes. Nature 2000; 404: 725–8.
- Hirsch J, Warkentin TE, Shaughnessy SG, Anand SS, Halperin JL, et al. Heparin and low molecular weight heparin. Mechanism of action, pharmacokinetics, dosing, monitoring, efficacy, and safety. Chest 2001; 119: 64S–94S.
- Béguin S, Welzel D, Al Dieri R, Hemker HC. Conjectures and refutations on the mode of action of heparins. The limited importance of anti-factor Xa activity as a pharmaceutical mechanism and a yardstick for therapy. Haemostasis 1999; 29: 170–8.
- Weitz JL. Low molecular-weight heparins. N Engl J Med 1997; 337: 688–98.
- Alban S. Pharmakologie der Heparine und der direkten Antikoagulantien. Hämostaseologie 2008; 5: 400–20.
- Hemker HC, Béguin S. Standard and method independent units for heparin anticoagulant activities. Thromb Haemost 1993; 70: 124–8.
- Toyoshima M, Nakajima M. Human heparanase. Purification, characterization, cloning and expression. J Biol Chem 1999; 274: 24153–60.
- Dong J, Kukula AK, Toyoshima M, Nakajima M. Genomic organization and chromosome localization of the newly identified heparanase gene. Gene 2000; 253: 171–8.
- Parish CR, Freeman C, Hulett MD. Heparanase: a key enzyme involved in cell invasion. Biochem Biophys Acta 2001; 1471: M99–M108.
- Hoogewerf AJ, Leone JW, Reardon IM, Howe WJ, Asa D, et al. CXCL chemokines connective tissue activating peptide-III and neutrophil activating peptide-2 are heparin/heparan sulfate-degrading enzymes. J Biol Chem 1995; 270: 3268–77.
- McKenzie E. Cloning and expression profiling of Hpa 2, a novel mammalian heparanase family member. Biochem Biophys Res Commun 2000; 276: 1170–7.
- Simizu S, Ishida K, Osada H. Heparanase as a molecular target of cancer chemotherapy. Cancer Sci 2004; 95: 553–8.
- Rohloff J, Zinke J, Schoppmeyer K, Tannapfel A, Witzigmann H, et al. Heparanase expression is a prognostic indicator for post-operative survival in pancreatic adenocarcinoma. Br J Cancer 2002; 86: 1270–5.
- Goldshmidt O, Zoharia E, Abramovitch R, Metzger S, Aingorn H, et al. Cell surface expression and secretion of heparanase markedly promote tumor angiogenesis and metastasis. Proc Natl Acad Sci USA 2002; 99: 10031–6.
- Linhardt RJ, Turnbull JE, Wang HM, Loganathan D, Gallagher JT. Examination of the substrate specificity of heparin and heparin sulphate lyases. Biochemistry 1990; 29: 2611–7.
- Desai UR, Wang HM, Linhardt RJ. Specificity studies on the heparin lyases from Flavobacterium heparinum. Biochemistry 1993; 32: 8140–5.
- Sie P, Ofosu F, Fernandez F, Buchanan MR, Petitou M, Boneu B. Respective role of antithrombin III and heparin cofactor II in the in vitro anticoagulant effect of heparin and various sulphated polysaccharides. Br J Haematol 1986; 64: 707–14.
- Gettins PG. Serpin structure, mechanism, and function. Chem Rev 2002; 102: 4751–804.
- Baglin TP, Carell RW, Church FC, Esmon CT, Huntington JA. Crystal structures of native and thrombin-complexed heparin cofactor II reveal a multistep allosteric mechanism. Proc Natl Acad Sci USA 2002; 99: 11079–84.
- Capila I, Linhardt RJ. Heparin-protein interactions. Angew Chem Int Ed 2002; 41: 390–412.
- Bourin MC, Lindahl U. Glycosaminoglycans and the regulation of blood coagulation. Biochem J 1999; 289: 313–30.
- Rezaie AR. Identification of basic residues in the heparin-binding exosite of factor Xa critical for heparin and factor Va binding. J Biol Chem 2000; 275: 3320–7.
- Schedin-Weiss S, Arocas V, Bock SC, Olson ST, Björk I. Specificity of the basic side chains of Lys114, Lys125 and Arg129 of antithrombin in heparin binding. Biochemistry 2002; 41: 12369–76.
- Thunberg L, Lindahl U, Tengblad A, Laurent TC, Jackson CM. On the molecular-weight dependence of anticoagulant activity of heparin. Biochem J 1979; 181: 241–3.
- Olson ST, Björk I, Sheffer R, Craig PA, Shore JD, Choay J. Role of the antithrombin-binding pentasaccharide in heparin acceleration of antithrombin-proteinase reactions. Resolution of the antithrombin conformational change contribution to heparin rate enhancement. J Biol Chem 1992; 267: 12528–38.

77. Broze GJ. Tissue factor pathway inhibitor. *Thromb Haemost* 1995; 74: 90–3.
78. Bajaj MS, Kuppuswamy MN, Saito H, Spitzer SG, Bajaj SP. Cultured normal human hepatocytes do not synthesise lipoprotein-associated coagulation inhibitor: evidence that epithelium is the principal site of synthesis. *Proc Natl Acad Sci* 1990; 87: 8869–73.
79. Hoppenstedt DA, Jeske W, Farsed J, Bermes EW jr. The role of tissue factor pathway inhibitor in the mediation of the antithrombotic actions of heparin and low molecular-weight heparin. *Blood Coagul Fibrinolysis* 1995; 6: S57–S64.
80. Cella G, Sbarai A, Mazzaro G, Motta G, Carraro P, et al. Tissue factor pathway inhibitor release induced by defibrinolytic and heparins. *Clin Appl Thromb Hemost* 2001; 7: 225–8.
81. Sandset PM, Bendz B, Hansen JB. Physiological function of tissue factor pathway inhibitor and interaction with heparins. *Haemostasis* 2000; 30 (Suppl 2): 48–56.
82. Warkentin TE. Heparin-induced thrombocytopenia: pathogenesis and management. *Br J Haematol* 2003; 121: 535–55.
83. Savi P, Pflieger AM, Héault JP, Michaux C, Duchaussoy P, et al. Comparative effects of two synthetic oligosaccharides on platelet activation induced by plasma from HIT patients. *J Thromb Haemost* 2003; 1: 2008–13.
84. Warkentin TE. An overview of heparin-induced thrombocytopenia syndrome. *Semin Thromb Hemost* 2004; 30: 273–83.
85. Greinacher A, Michels I, Mueller-Eckhardt C. Heparin-associated thrombocytopenia: the antibody is not heparin-specific. *Thromb Haemost* 1992; 67: 545–9.
86. Raoou I, Zhai L, Kowalska MA, Arepally GM, Cines DB, Poncz M. Ultralarge complexes of PF4 and heparin are central to the pathogenesis of heparin-induced thrombocytopenia. *Blood* 2005; 105: 131–8.
87. Würfel W. Habituelle Aborte. Diagnostik und Therapie der Thrombophilie. *Gynäkol Prax* 2010; 34: 427–36.
88. Greinacher A. Heparininduzierte Thrombozytopenie. *Dtsch Arztebl* 2003; 100: A2220–9.
89. Harenberg J, Heilmann L, Rath W, Riess H, von Tempelhoff GF, et al. Niedermolekulare Heparine in der Schwangerschaft. *Dtsch Arztebl* 2002; 99: A424–34.
90. Hardtke K, Böhme H (Hrsg). Kommentar zum Europäischen Arzneibuch Ph. Eur. 5.0, Protaminsulfat. Wissenschaftliche Verlagsgesellschaft, Stuttgart 2005.
91. Hirsh J, Warkentin TE, Raschke R, Granger C, Ohman EM, Dalen JE. Heparin and low molecular weight heparin: mechanisms of action, pharmacokinetics, dosing considerations, monitoring, efficacy, and safety. *Chest* 1999; 115: S489–510.
92. Lüllmann H, Mohr K, Hein L. *Pharmakologie und Toxikologie*. 17. Aufl. Georg-Thieme-Verlag, Stuttgart New York, 2010.
93. Hänsel R, Sticker O (Hrsg). *Pharmakognosie – Phytopharmazie*. 9. Aufl. Springer Medizin Verlag, Heidelberg, 2010.
94. Lemmer B, Brune K (Hrsg). *Klinische Pharmakologie*. 14. Aufl. Springer Medizin Verlag, Heidelberg, 2010.
95. HajMohammadi S, Enjiyoji K, Princivalle P, Lech M, Beeler D, et al. Normal levels of anticoagulant heparin sulphate are not essential for normal hemostasis. *J Clin Invest* 2003; 111: 989–99.
96. Alban S. *Pharmakologie der Heparine und der direkten Antikoagulanzen*. Hämostaseologie 2009; 5: 400–20.
97. Rasche H. *Niedermolekulare Heparine*. *Dtsch Arztebl* 1988; 41: A1007–13.
98. Rabenstein DL. Heparin and heparan sulfate: structure and function. *Nat Prod Rep* 2002; 19: 312–31.
99. Jaques LB, Debnath AK. Simultaneous evaluation of tissue heparin and mast cells in small tissue samples. *Am J Physiol* 1970; 219: 1155–60.
100. Müller-Eberhard HJ. Molecular organization and function of the complement system. *Annu Rev Biochem* 1988; 57: 321–47.
101. Bohana-Kashtan O, Ziporen L, Donin N, Kraus S, Fishelson Z. Cell signals transduced by complement. *Mol Immunol* 2004; 41: 583–97.
102. Makrides SC. Therapeutic inhibition of the complement system. *Pharmacol Rev* 1998; 50: 59–88.
103. Edens RE, Linhardt RJ, Weiler JM. Heparin is not just an anticoagulant anymore: six and one-half decades of studies on the ability of heparin to regulate complement activity. *Complement profiles* 1993; 1: 96–120.
104. Pangburn MK. Analysis of recognition in the alternative pathway of complement. Effect of polysaccharide size. *J Immunol* 1989; 142: 2766–70.
105. Strunk R, Colten HR. Inhibition of the enzymatic activity of the first component (C1) by heparin. *Clin Immunol Immunopath* 1976; 6: 248–55.
106. Weiler JM, Yurt RW, Fearon DT, Austen F. Modulation of the formation of the amplification convertase of complement, C3b, Bb, by native and commercial heparin. *J Exp med* 1978; 147: 409–21.
107. Sahu A, Pangburn MK. Identification of multiple sites of interaction between heparin and the complement system. *Mol Immunol* 1993; 30: 679–84.
108. Yu H, Munoz EM, Edens RE, Linhardt RJ. Kinetic studies on the interaction of heparin and complement proteins using Plasmon surface resonance. *Biochim Biophys Acta* 2005; 1726: 168–76.
109. Ecker EE, Gross P. Anticomplementary power of heparin. *J Infect Dis* 1929; 44: 250–3.
110. Itoh K, Nakao A, Kishimoto W, Takagi H. Heparin effects on superoxide production by neutrophils. *Eur Surg Res* 1995; 27: 184–8.
111. Dandona P, Qutob T, Hamouda W, Bakri F, Aljada A, Kumbkarni Y. Heparin inhibits reactive oxygen species generation by polymorphonuclear cells and mononuclear leucocytes. *Thromb Res* 1999; 96: 437–43.
112. Attanasio M, Gori AM, Giusti B, Pepe G, Ceglie P, et al. Cytokine gene expression in human LPS- and IFN-gamma stimulated mononuclear cells is inhibited by heparin. *Thromb Haemost* 1998; 79: 959–62.
113. Lantz M, Thysell H, Nilsson E, Olsson I. On the binding of tumor necrosis factor (TNF) to heparin and the release in vitro of the TNF-binding protein I by heparin. *J Clin Invest* 1991; 88: 2026–31.
114. Meyer J, Cox CS, Herndon DN, Nakazawa H, Lentz CW, et al. Heparin in experimental hyperdynamic sepsis. *Crit Care Med* 1993; 21: 84–9.
115. Vancheri C, Mastruzzo C, Armato F, Tomaselli V, Magri S, et al. Intranasal heparin reduces eosinophil recruitment after nasal allergen challenge in patients with allergic rhinitis. *J Allergy Clin Immunol* 2001; 108: 703–8.
116. Gaffney A, Gaffney P. Rheumatoid arthritis and heparin. *Br J Rheumatol* 1996; 35: 808–10.
117. Bowler SD, Smith SM, Lavercombe PS. Heparin inhibits the immediate response to antigen in the skin and lungs of allergic subjects. *Am Rev Respir Dis* 1993; 147: 160–3.
118. Tyrrell DJ, Home AP, Holme KR, Preuss JM, Page CP. Heparin in inflammation: potential therapeutic applications beyond anticoagulation. *Adv Pharmacol* 1999; 46: 151–208.
119. Nassar RA, Browne EP, Chen J, Klibanov A. Removing human immunodeficiency virus (HIV) from human blood using immobilized heparin. *Biotechnol Lett* 2012; 34: 853–6.
120. Nahmias AJ, Kibrick S. Inhibitory effect of heparin on herpes simplex virus. *J Bacteriol* 194; 87: 1060–6.
121. Lin YL, Lei HY, Lin YS, Yeh TM, Chen SH, Liu HS. Heparin inhibits dengue-2 virus infection of five human liver cell lines. *Antiviral Res* 2002; 56: 93–6.
122. Dennis LH, Conrad ME. Anticoagulant and antimalarial action of heparin in simian malaria. *The Lancet* 1968; 291: 769–71.
123. Boyle MJ, Richards JS, Gilson PR, Chai W, Beeson JG. Interactions with heparin-like molecules during erythrocyte invasion by *Plasmodium falciparum* merozoites. *Blood* 2010; 115: 4559–68.
124. De Witte L, Bobardt M, Chatterji U, Degeest G, David G, Geijtenbeek TB, Gallay P. Syndecan-3 is a dendritic cell-specific attachment receptor for HIV-1. *PNAS* 2007; 104: 19464–9.
125. Salvador B, Sexton NR, Carrion R jr, Nunneley J, Patterson JL, et al. Filoviruses utilize glycosaminoglycans for their attachment to target cells. *J Virol* 2013; 87: 3295–304.
126. Bobardt MD, Chatterji U, Selvarajah S, Van der Schueren B, David G, et al. Cell-free human immunodeficiency virus type 1 transcytosis through primary genital epithelial cells. *J Virol* 2007; 81: 395–405.
127. Metz M. *Physiologische Funktionen der Mastzellen*. Habilitationsschrift, Charité-Universitätsmedizin Berlin, 2009.
128. Xia CQ, Kao KJ. Heparin induces differentiation of CD1a+ dendritic cells from monocytes: phenotypic and functional characterization. *J Immunol* 2002; 168: 1131–8.
129. Shriver Z, Liu D, Sasisekharan R. Emerging views of heparan sulfate glycosaminoglycan structure-activity relationships modulating dynamic biological functions. *Trends Cardiovasc Med* 2002; 12: 71–7.
130. Ornitz DM, Itoh N. Fibroblast growth factors. *Genome Biol* 2001; 2: R3005.
131. Cross MJ, Claesson-Welsh L. FGF and VEGF function in angiogenesis. Signaling pathways, biological responses and therapeutic inhibition. *Trends Pharmacol Sci* 2001; 22: 201–7.
132. Monsa SA, Kaiser B. Tissue factor pathway inhibitor in thrombosis and beyond: role of heparin. *Drugs Future* 2004; 29: 751–6.
133. Ibrahim OA, Zhang F, Hrstka SC, Mohammadi M, Linhardt RJ. Kinetic model for FGF, FGFR and proteoglycan signal transduction complex assembly. *Biochemistry* 2004; 43: 4724–30.
134. Pye DA, Vives RR, Turnbull JE, Hyde P, Gallagher JT. Heparan sulfate oligosaccharides require 6-O-sulfation for promotion of basic fibroblast growth factor mitogenic activity. *J Biol Chem* 1998; 273: 22936–42.
135. Giumond S, Maccarana M, Olwin BB, Lindahl U, Rapraeger AC. Activating and inhibitory heparin sequences for FGF-2 (basic FGF). Distinct requirements for FGF-1, FGF-2, and FGF-4. *J Biol Chem* 1993; 268: 23906–14.
136. Fannon M, Forsten KE, Nugent MA. Potentiation and inhibition of bFGF binding by heparin. A model for regulation of cellular response. *Biochemistry* 2000; 39: 1434–45.
137. Forsten KE, Fannon M, Nugent MA. Potential mechanisms for the regulation of growth factor binding by heparin. *J Theor Biol* 2000; 205: 215–30.
138. Parish CR, Freeman C, Hulett MD. Heparanase: a key enzyme involved in cell invasion. *Biochim Biophys Acta* 2001; 1471: M99–108.
139. Gong F, Jemth P, Escobar Galvis ML, Vlodavsky I, Horner A, Lindahl U, Li JP. Processing of macromolecular heparin by heparanase. *J Biol Chem* 2003; 278: 35152–8.
140. Castellot JJ jr, Wong K, Herman B, Hoover RL, Albertini DF, et al. Binding and internalization of heparin by vascular smooth muscle cells. *J Cell Physiol* 1985; 124: 13–20.
141. Letourneur D, Caleb BL, Castellot JJ jr. Heparin binding, internalization, and metabolism in vascular smooth muscle cells. I. Upregulation of heparin binding correlates with antiproliferative activity. *J Cell Physiol* 1995; 165: 676–86.
142. Letourneur D, Caleb BL, Castellot JJ jr. Heparin binding, internalization, and metabolism in vascular smooth muscle cells. II. Degradation and secretion in sensitive and resistant cells. *J Cell Physiol* 1995; 165: 687–95.
143. Patton WA 2nd, Granzow CA, Getts LA, Thomas SC, Zotter LM, et al. Identification of a heparin-binding protein using monoclonal antibodies that block heparin binding to porcine aortic endothelial cells. *Biochem J* 1995; 311 (III): 461–9.
144. Delmolino LM, Castellot JJ jr. Heparin suppresses sgk, an early response gene in proliferating vascular smooth muscle cells. *J Cell Physiol* 1997; 178: 371–9.
145. Mishra-Gorur K, Castellot JJ jr. Heparin rapidly and selectively regulates tyrosine phosphorylation in vascular smooth muscle cells. *J Cell Physiol* 1999; 178: 205–15.
146. Daum G, Hedin U, Wang Y, Wang T, Clowes AW. Diverse effects of heparin on mitogen-activated protein kinase dependent signal transduction in vascular smooth muscle cells. *Circ Res* 1997; 81: 17–23.
147. Mishra-Gorur K, Singer HA, Castellot JJ jr. Heparin inhibits phosphorylation and autonomous activity of Ca(2+)/calmodulin-dependent protein kinase II in vascular smooth muscle cells. *Am J Pathol* 2002; 161: 1893–901.
148. Castellot JJ jr, Beeler DL, Rosenberg RD, Karnovsky MJ. Structural determinants of the capacity of heparin to inhibit the proliferation of vascular smooth muscle cells. *J Cell Physiol* 1984; 120: 315–20.
149. Wright TC, Castellot JJ jr, Petitou M, Lormeau JC, Choay J, Karnovsky MJ. Structural determinants of heparin's growth inhibitory activity. Interdependence of oligosaccharide size and charge. *J Biol Chem* 1989; 264: 1534–42.
150. Faham S, Linhardt RJ, Rees DC. Diversity does make a difference: fibroblast growth factor-heparin interactions. *Curr Opin Struct Biol* 2001; 11: 629–34.
151. De Agostini AI. An unexpected role for anticoagulant heparan sulfate proteoglycans in reproduction. *Swiss Med Wkly* 2006; 136: 583–90.
152. Liu D, Shriver Z, Qi Y, Venkataraman G, Sasisekharan. Dynamic regulation of tumor growth and metastasis by heparin sulfate glycosaminoglycans. *Semin Thromb Hemost* 2002; 28: 67–78.
153. Castelli R, Porro F, Tarsia P. The heparins and cancer: review of clinical trials and biological properties. *Vasc Med* 2004; 9: 2005–13.
154. Sanderson RD. Heparan sulfate proteoglycans in invasion and metastasis. *Semin Cell Dev Biol* 2001; 12: 89–98.
155. Filmus J. Glypicans in growth control and cancer. *Glycobiology* 2001; 11: 19R–23R.
156. Jiang X, Couchman JR. Perlecan and tumor angiogenesis. *J Histochem Cytochem* 2003; 51: 1393–410.

157. Bernfield M, Kokenyesi R, Kato M, Hinkes MT, Spring J, Gallo RL, Loe E. Biology of the syndecans: a family of transmembrane heparin sulfate proteoglycans. *Annu Rev Cell Biol* 1992; 8: 365–93.
158. Lander AD, Selleck SB. The elusive functions of proteoglycans: in vivo veritas. *J Cell Biol* 2000; 148: 237–42.
159. Liu X, McFarland DC, Nestor KE, Velleman SG. Developmental regulated expression of syndecan-1 and glypican in pectoralis major muscle in turkeys with different growth rates. *Develop Growth Differ* 2004; 46: 37–51.
160. David G. Integral membrane heparin sulfate proteoglycans. *FASEB J* 1993; 7: 1023–30.
161. Kim CW, Golberger OA, Gallo RL, Bernfield M. Members of the syndecan family of heparin sulfate proteoglycans are expressed in distinct cell-, tissue-, and developmental-specific patterns. *Mol Biol Cell* 1994; 5: 797–805.
162. Baciuc PC, Acaster C, Goetinick PF. Molecular cloning and genomic organization of chicken syndecan-4. *J Biol Chem* 1994; 269: 696–703.
163. Chen I, KlassC, Woods A. Syndecan-2 regulates transforming factor- $\beta$  signaling. *J Biol Chem* 2004; 279: 15715–8.
164. Woods A, Couchman JR. Syndecan-4 heparan sulfate proteoglycan is a selectively enriched and widespread focal adhesion component. *Mol Biol Cell* 1994; 5: 183–92.
165. Rapraeger AC. Molecular interactions of syndecan during development. *Cell Dev Biol* 2001; 12: 107–16.
166. Sutherland AE, Sanderson RD, Mayes M, Seibert M, Calarco PG, et al. Expression of syndecan, a putative low affinity growth factor, in the early mouse embryo. *Development* 1991; 113: 339–51.
167. Trautman MS, Kimelman J, Bernfield M. Developmental expression of syndecan, an integral membrane proteoglycan, correlates with cell differentiation. *Development* 1989; 111: 213–20.
168. Yamaguchi Y. Heparan sulfate proteoglycans in the nervous system: their diverse roles in neurogenesis, axon guidance, and synaptogenesis. *Sem Cell Dev Biol* 2001; 12: 99–106.
169. Fuentaltba L, Carey DJ, Brandon E. Antisense inhibition of syndecan-3 expression during skeletal muscle differentiation accelerates myogenesis through a basic fibroblast growth factor-dependant mechanism. *J Biol Chem* 1999; 274: 37878–84.
170. Cornelison DD, Filla MS, Stanley HM, Rapraeger AC, Olwin BB. Syndecan-3 and syndecan-4 specifically mark skeletal muscle satellite cells and are implicated in satellite cell maintenance and muscle regeneration. *Dev Biol* 2001; 239: 79–94.
171. Casar JC, Cabello-Verrugio C, Olguin H, Aldunate R, Inestrosa NC, Brandon E. Heparan sulfate proteoglycans are increased during skeletal muscle regeneration: requirement of syndecan-3 for successful fiber formation. *J Cell Sci* 2004; 117: 73–84.
172. Mertens G, Van der Schueren B, van den Bergh E, David G. Heparan sulphate expression in polarized epithelial cells: the apical sorting of glypican (GPI-anchored proteoglycan) is inversely related to its heparin sulphate content. *J Cell Biol* 1996; 132: 487–97.
173. Liu X, McFarland DC, Nestor KE, Velleman SG. Developmental regulated expression of syndecan-1 and glypican in pectoralis major muscle in turkeys with different growth rates. *Develop Growth Differ* 2004; 46: 37–51.
174. Stipp CS, Litwack ED, Lander AD. Cerebroglycan: an integral membrane heparin sulfate proteoglycan that is unique to the developing nervous system and expressed specifically during neuronal differentiation. *J Cell Biol* 1994; 124: 149–60.
175. Pellegrini M, Pilia G, Pantano S, Lucchini F, Uda M, et al. Gpc3 expression correlates with the phenotype of the Simpson-Golabi-Beihmel syndrome. *Dev Dyn* 1998; 213: 431–9.
176. Watanabe K, Yamada H, Yamaguchi Y. K-glypican: a novel GPI-anchored heparin sulphate proteoglycan that is highly expressed in developing brain and kidney. *J Cell Biol* 1995; 130: 1207–18.
177. Saunders S, Paine-Saunders S, Lander AD. Expression of the cell surface proteoglycan glypican-5 is developmentally regulated in kidney, limb and brain. *Dev Biol* 1997; 190: 78–93.
178. Veuglers M, De Cat B, Ceulemans H, Bruystens AM, Coomans C, et al. Glypican-6, a new member of the glypican family of cell surface heparin sulphate proteoglycans. *J Biol Chem* 1999; 38: 26968–77.
179. Gonzalez AD, Kaya M, Shi W, Song H, Testa JR, et al. OCL-5/GPC3, a glypican encoded by a gene that is mutated in the Simpson-Golabi-Beihmel overgrowth syndrome, induces apoptosis in a cell line specific manner. *J Cell Biol* 1998; 141: 1407–14.
180. Pilia G, Hughes-Benzie RM, MacKenzie A, Baybayan P, Chen EY, et al. Mutations in GPC3, a glypican gene, cause the Simpson-Golabi-Beihmel overgrowth syndrome. *Nat Genet* 1996; 12: 241–7.
181. Veuglers M, Cat BD, Muijldermans SY, Reekmans G, Delande N, et al. Mutational analysis of GPC3/GPC4 glypican gene cluster on Xq26 in patients with Simpson-Golabi-Beihmel overgrowth syndrome: identification of loss-of-function mutations in the GPC3 gene. *Hum Mol Genet* 2000; 9: 1321–8.
182. Qian X, Davis AD, Goderle SK, Temple S. FGF2 concentration regulates the generation of neurons and glia from multipotent cortical stem cells. *Cell Biol* 1997; 18: 81–93.
183. Hagihara K, Watanabe K, Chun J, Yamaguchi Y. Glypican-4 is an FGF2-binding heparin sulphate proteoglycan expressed in neural precursor cells. *Dev Dyn* 2000; 219: 353–67.
184. Cheifetz S, Bellón T, Calés C, Vera S, Bernbeu C, et al. Endoglin is a component of the transforming growth factor-beta receptor system in human endothelial cells. *J Biol Chem* 1992; 267: 19027–30.
185. Lopez-Casillas F, Wrana JL, Massague J. Betaglycan presents ligands to the TGF beta signaling receptor. *Cell* 1993; 73: 1435–44.
186. Blobel GC, Liu X, Fang SJ, How T, Lodish HF. A novel mechanism for regulating transforming growth factor interaction with the PDZ domain protein, GIPC. *J Biol Chem* 2001; 43: 39608–17.
187. Velasco-Loyden G, Arribas J, Lopez-Casillas F. The shedding of betaglycan is regulated by pervanadate and mediated by membrane type matrix metalloproteinase-1. *J Biol Chem* 2004; 279: 7721–33.
188. Screaton GR, Bell MV, Jackson DG, Cornelis FB, Gerth U, Bell JI. Genomic structure of DNA encoding the lymphocyte homing receptor CD44 reveals at least 12 alternatively spliced exons. *Proc Natl Acad Sci USA* 1992; 89: 12160–4.
189. Goodison S, Urquidí V, Tarin D. CD44 cell adhesion molecules. *Mol Path* 1999; 52: 189–96.
190. Tuhkanen AL, Tammi M, Tammi R. CD44 substituted with heparan sulfate and endo-beta-galactosidase-sensitive oligosaccharides: a major glycoprotein in adult human epidermis. *J Invest Dermatol* 1997; 109: 213–8.
191. Sackstein R, Merzaban JS, Cain DW, Dagia NM, Spencer JA, Lin CP, Wohlgemuth R. Ex vivo glycan engineering of CD44 programs human multipotent mesenchymal stromal cell trafficking to bone. *Nat Med* 2008; 14: 181–7.
192. Napier SL, Healy ZR, Schnaar RL, Konstantopoulos K. Selectin ligand expression regulates the initial vascular interactions of colon carcinoma cells: the roles of CD44v and alternative sialofucosylated selectin ligands. *J Biol Chem* 2007; 282: 3433–41.
193. Oxley SM, Sackstein R. Detection of an L-selectin ligand on hemopoietic progenitor cell line. *Blood* 1994; 84: 3299–306.
194. Griffith JS, Liu YG, Tekmal RR, Binkley PA, Holden AEC, Schenken RS. Menstrual endometrial cells from women with endometriosis demonstrate increased adherence to peritoneal cells and increased expression of CD44 splice. *Fertil Steril* 2010; 93: 1745–9.
195. Hassell JR, Robey PG, Barrach HJ, Wilczek J, Rennard SI, Martin GR. Isolation of a heparin sulfate-containing proteoglycan from basement membrane. *Proc Natl Acad Sci USA* 1980; 77: 4494–8.
196. Ledbetter SR, Fisher LW, Hassell JR. Domain structure of the basement membrane heparan sulfate proteoglycan. *Biochemistry* 1987; 26: 988–95.
197. Hopf M, Göhring W, Kofeldt E, Yamada Y, Timpl R. Recombinant domain IV of perlecan binds to nidogens, laminin-nidogen complex, fibronectin, fibulin-2 and heparin. *Eur J Biochem* 1999; 259: 917–25.
198. Brown JC, Sasaki T, Göhring W, Yamada Y, Timpl R. The C-terminal domain V of perlecan promotes beta1 integrin-mediated cell adhesion, binds heparin, nidogen and fibulin-2 and can be modified by glycosaminoglycans. *Eur J Biochem* 1997; 250: 39–46.
199. Olson BR. Life without perlecan has its problems. *J Cell Biol* 1999; 147: 90–911.
200. Arikawa-Hirshawa E, Watanabe H, Takami H, Hassell JR, Yamada Y. Perlecan is essential for cartilage and cephalic development. *Nat Genet* 1999; 23: 354–8.
201. Costell M, Gustafsson E, Aszódi A, Mörgelin M, Bloch W, et al. Perlecan maintains integrity of cartilage and some basement membranes. *J Cell Biol* 1999; 147: 1109–22.
202. Bayne EK, Anderson MJ, Farnborough DM. Extracellular matrix organization in developing muscle: correlation with acetylcholine receptor aggregates. *J Cell Biol* 1984; 99: 1486–501.
203. Hassell JR, Yamada Y, Arikawa-Hirasawa E. Role of perlecan in skeletal development and diseases. *Glycoconjugate J* 2003; 19: 263–7.
204. Arikawa-Hirasawa E, Rossi SG, Rotundo RL, Yamada Y. Absence of acetylcholinesterase at the neuromuscular junctions of perlecan-null mice. *Nat Neurosci* 2002; 5: 119–23.
205. Arikawa-Hirasawa E, Le AH, Nishino I, Nonaka I, Ho NC, et al. Structural and functional mutations of the perlecan gene cause Schwartz-Jampel syndrome, with myotonic myopathy and chondrodysplasia. *A J Hum Genet* 2002; 70: 1368–75.
206. Arikawa-Hirasawa E, Wilcox WR, Le AH, Silverman N, Govindraj P, et al. Dyssegmental dysplasia, Silverman-Handmaker type, is caused by functional null mutations of the perlecan gene. *Nat Genet* 2001; 27: 431–4.
207. Stetefeld J, Alexandrescu AT, Maciejewski MW, Jenny M, Rathgeb-Szabo K, et al. Modulation of agrin function by alternative splicing and Ca<sup>2+</sup> binding. *Structure* 2004; 12: 503–15.
208. Kröger S, Schröder JE. Agrin in the developing CNS: new roles for a synapse organizer. *News Physiol Sci* 2002; 17: 202–17.
209. Groffen AJ, Buskens CA, van Kuppevelt TH, Veerkamp JH, Monnens LA, van den Heuvel LP. Primary structure and high expression of human agrin in basement membranes of adult lung and kidney. *Eur J Biochem* 1998; 254: 123–8.
210. Mantych KB, Ferreira A. Agrin differentially regulates the rates of axonal and dendritic elongation in cultured hippocampal neurons. *J Neurosci* 2001; 21: 6802–9.
211. Gautam M, Noakes PG, Moscoso L, Rupp F, Scheller RH, Merlie JP, Sanes JR. Defective neuromuscular synaptogenesis in agrin-deficient mutant mice. *Cell* 1996; 85: 525–35.
212. Sanes JR, Apel ED, Gautam M, Glass D, Grady RM, et al. Agrin receptors at the skeletal neuromuscular junction. *Ann NY Acad Sci* 1998; 841: 1–13.
213. Franke CW, Bruckner P, Bruckner-Tuderman L. Collagenous transmembrane proteins: recent insights into biology and pathology. *J Biol Chem* 2005; 280: 4005–8.
214. Dong S, Cole GJ, Halfter W. Expression of collagen XVIII and localization of its glycosaminoglycan attachment sites. *J Biol Chem* 2003; 278: 1700–7.
215. O'Reilly MS, Boehme T, Shing Y, Fukai N, Vasios G, et al. Endostatin: an endogenous inhibitor of angiogenesis and tumor growth. *Cell* 1997; 88: 277–85.
216. Benezra R, Rafii S. Endostatin's endpoints-deciphering the endostatin antiangiogenic pathway. *Cancer Cell* 2004; 5: 2005–6.
217. Jing Y, Lu H, Wu K, Subramanian IV, Ramakrishnan S. Inhibition of ovarian cancer by RGD-P125 A-endostatin-Fc fusion proteins. *Int J Cancer* 2011; 129: 751–6.
218. Huang G, Chen L. Recombinant human endostatin improves anti-tumor efficacy of paclitaxel by normalizing tumor vasculature in Lewis lung carcinoma. *J Cancer Res Clin Oncol* 2010; 136: 1201–11.

# Mitteilungen aus der Redaktion

Besuchen Sie unsere Rubrik

## ☒ Medizintechnik-Produkte



Neues CRT-D Implantat  
Intica 7 HF-T QP von Biotronik



Artis pheno  
Siemens Healthcare Diagnostics GmbH



Philips Azurion:  
Innovative Bildgebungslösung

Aspirator 3  
Labotect GmbH



InControl 1050  
Labotect GmbH

## e-Journal-Abo

Beziehen Sie die elektronischen Ausgaben dieser Zeitschrift hier.

Die Lieferung umfasst 4–5 Ausgaben pro Jahr zzgl. allfälliger Sonderhefte.

Unsere e-Journale stehen als PDF-Datei zur Verfügung und sind auf den meisten der marktüblichen e-Book-Readern, Tablets sowie auf iPad funktionsfähig.

## ☒ Bestellung e-Journal-Abo

### Haftungsausschluss

Die in unseren Webseiten publizierten Informationen richten sich **ausschließlich an geprüfte und autorisierte medizinische Berufsgruppen** und entbinden nicht von der ärztlichen Sorgfaltspflicht sowie von einer ausführlichen Patientenaufklärung über therapeutische Optionen und deren Wirkungen bzw. Nebenwirkungen. Die entsprechenden Angaben werden von den Autoren mit der größten Sorgfalt recherchiert und zusammengestellt. Die angegebenen Dosierungen sind im Einzelfall anhand der Fachinformationen zu überprüfen. Weder die Autoren, noch die tragenden Gesellschaften noch der Verlag übernehmen irgendwelche Haftungsansprüche.

Bitte beachten Sie auch diese Seiten:

**Impressum**

**Disclaimers & Copyright**

**Datenschutzerklärung**