

Journal für

# Reproduktionsmedizin und Endokrinologie

– Journal of Reproductive Medicine and Endocrinology –

Andrologie • Embryologie & Biologie • Endokrinologie • Ethik & Recht • Genetik  
Gynäkologie • Kontrazeption • Psychosomatik • Reproduktionsmedizin • Urologie



## Estrogen Free Contraception: Progestin-only-systems

Ahrendt HJ, Bühling KJ

*J. Reproduktionsmed. Endokrinol* 2010; 7 (Sonderheft  
1), 97-105

[www.kup.at/repromedizin](http://www.kup.at/repromedizin)

Online-Datenbank mit Autoren- und Stichwortsuche

Offizielles Organ: AGRBM, BRZ, DVR, DGA, DGGEF, DGRM, D-I-R, EFA, OEGRM, SRBM/DGE

Indexed in EMBASE/Excerpta Medica/Scopus

Krause & Pachernegg GmbH, Verlag für Medizin und Wirtschaft, A-3003 Gablitz

# Estrogen Free Contraception: Progestin-only-systems\*

H.-J. Ahrendt<sup>1</sup>, K. J. Böhling<sup>2</sup>

To reduce side effects of estrogen-progestin combination preparations, the dose of estrogen has continuously been reduced in the pill. As an alternative, estrogen-free preparations are increasingly used and are now available as oral, subdermal, intrauterine and intramuscular applications. The benefits of estrogen-free contraceptives are the prevention of estrogen-related side effects (nausea, edema, weight gain, mastodynia) and of cycle-dependent side effects (dysmenorrhea, pelvic pain, premenstrual syndrome [PMS], hypermenorrhea, menstrual migraine). Furthermore, they can be used in women with risk factors, in whom estrogens are contraindicated. These include hypertension, thrombophilia, status post thrombosis, myocardial infarction, stroke, liver tumors, cholelithiasis and during lactation. In the following sections, the available preparations are being discussed. **J Reproduktionsmed Endokrinol 2010; 7 (Special Issue 1): 97–105.**

**Key words:** estrogen free contraception, estrogen free pills, implants, intrauterine hormonal system, injections

## ■ Introduction

Apart from combination preparations, which contain estrogens and progestins, various estrogen-free systems are now available in different applications (oral, subdermal, intrauterine, intramuscular).

The reduction in ethinyl estradiol dosage from an initial 75 µg to 50 µg and further down to 20 µg and 15 µg in combination preparations (combined oral contraceptives, COCs), the introduction of estradiol valerate in the COCs and particularly the absence of estrogens in hormonal contraceptives are recent developments that are thought to avoid estrogen-dependent side effects and, in particular, reduce the risk for thrombosis [1, 2].

The dosage regimen of combination preparations is usually 21 to 26 days of active therapy, followed by a 2–7-day interval in order to induce withdrawal bleeding.

However, progestin contraceptives (progestin-only systems) are taken continuously without pill-free intervals and progestin-only devices continuously release equal hormone doses. This, in turn, avoids hormonal fluctuation and cycle-dependent complaints.

The progestins greatly account for the high contraceptive efficacy of combined hormonal contraceptives. Important mechanisms of action are the central inhibition of the hypothalamic-pituitary-

ovarian axis (inhibition of ovulation) as well as the inhibition of the three peripheral parameters cervix, endometrium and tubal function (Tab. 1).

The following estrogen-free preparations are currently available:

- Oral progestin preparations (Progestin-only-Pills, POPs)
  - incomplete ovulation inhibition: 28 mini<sup>®</sup>, Microlut<sup>®</sup>, Micro-30 Wyeth<sup>®</sup>, containing 0.03 mg levonorgestrel (LNG) each complete ovulation inhibition: Cerazette<sup>®</sup> containing 0.075 mg desogestrel
- Depot injections
  - DepoClinovir<sup>®</sup> with 150 mg medroxyprogesterone acetate (MPA)
  - Noristerat<sup>®</sup> containing 200 mg norethisterone enanthate
- Subdermal implants: Implanon<sup>®</sup>, containing etonogestrel
- Intrauterine devices: Mirena<sup>®</sup> containing levonorgestrel
- Postcoital contraception containing levonorgestrel (Unofem<sup>®</sup> and Levonon<sup>®</sup>) and Ulipristal (ellaOne<sup>®</sup>)

## ■ Advantages of Progestin-only Preparations

Referring to Ahrendt [3] and Rabe et al. [4], the progestin-only preparations have the following advantages:

- No estrogen-dependent side effects
  - Nausea
  - Edema
  - Weight gain
  - Mastodynia
  - Estrogen-dependent headache
- No cycle-dependent complaints
  - Dysmenorrhea
  - Pelvic pain, premenstrual syndrome (PMS)
  - Hypermenorrhea
  - Menstrual migraine
- Compared to combined oral contraceptives reduced risk for
  - Stroke
  - Myocardial infarction
  - Thrombosis
- Reduced risk for endometrial cancer
- Applicable in patients with contraindications for combined estrogen-progestin preparations


**Table 1:** Mechanism of action of progestin only pills (POPs)

	Cervical mucus	Endometrium	Fallopian tube	Ovulation inhibition
Mini pill (0.03 mg levonorgestrel)	x	x	x	–
Estrogen-free pill (0.075 mg desogestrel)	x	x	x	x
Depot injections (medroxyprogesterone acetate, norethisterone enanthate)	x	x	x	x
Etonogestrel implant	x	x	x	x
Intrauterine devices (levonorgestrel)	x	x	x	–

\* Reprint of: Ahrendt HJ. Estrogenfreie Kontrazeption. Progestin-only-Systeme. Gynäkol Geburtsmed Gynäkol Endokrinol 2009; 5: 152–66. Reprinted with permission by akademos Wissenschaftsverlag GmbH, Hamburg. Translation: Ahrendt H.-J.

From the <sup>1</sup>Praxis für Frauenheilkunde, Klinische Forschung und Weiterbildung, Magdeburg, and the <sup>2</sup>Universitätsklinikum Hamburg-Eppendorf, Hamburg, Germany

**Correspondence:** Prof. Hans-Joachim Ahrendt, MD, Praxis für Frauenheilkunde, Klinische Forschung und Weiterbildung, Halberstädter Straße 122, D-39112 Magdeburg, Germany; e-mail: ahrendt@prof-ahrendt-frauenarzt.de



**Figure 1:** Change of estrogen-dependent side effects after a three month treatment with estrogen-free pills (0.75 mg Desogestrel), n = 406

- Hypertension (provided that treatment is initiated)
- Thrombophilia
- Status post thrombosis
- Myocardial infarction
- Stroke
- Liver tumors
- Cholelithiasis
- Suitable during lactation
- Further benefits of non-oral progestin-only systems (subdermal implant, intrauterine device, depot injections)
  - Bypassing of gastrointestinal tract and the first-pass effect
  - No continuous tablet intake, thus also suitable for patients with lack of compliance
  - High contraceptive efficacy
  - Safe use in patients with contraindications for estrogen-progestin combination preparations

### ■ Advantages of Estrogen-free Contraception

The most commonly listed estrogen-dependent side effects of estrogen-progestin combination preparations are mastodynia, fullness, nausea and mood swings as well as headaches during pill intake [5–7].

A multi-centric observational trial in Germany investigated the situation of 403 patients that presented with estrogen-dependent complaints such as nausea, mastodynia, headaches and edemas due to the intake of combined oral con-

traceptives. After changing to the continuous progestin intake of desogestrel 75 µg/day (Cerazette®), the estrogen-dependent side effects ameliorated significantly after three cycles. Nausea was the most frequent symptom (92 % of women), followed by mastodynia (90 % of women), headaches (84 % of women) and edemas (74 % of women) [8]. These results were confirmed by an observational study in the Czech Republic (n = 484) [9] (Fig. 1).

### ■ Elimination of Cycle-dependent Complaints

#### Primary dysmenorrhea

Menstrual pain or dysmenorrhea affects approximately 50 % of all women [10], which, in turn, has a great impact on the women's daily routine and their ability to attend school or work.

Menstruation-associated pain results from prostaglandin-induced uterine contractions and uterine ischemia [10–13]. Estrogen-progestin combination preparations are an effective therapy for primary dysmenorrhea, as they suppress ovulation and decrease endometrial thickness [10–12, 14]. However, as cyclic combination preparations induce withdrawal bleeding, some women remain with menstrual cramps. Results from a meta-analysis of six randomized, controlled studies showed that the effectiveness of combination preparations in dysmenorrhea increases, if the pill-free

interval is reduced [15]. By shortening the pill-free interval in the new combination preparations to four or two days, by using the pills in an extended-cycle dosage regimen or even with continuous intake, the further reduction of dysmenorrhea is feasible [16].

Progestin only systems such as progestin only pills (POPs), intrauterine devices (IUDs), implants and depot injections, which have no scheduled pill-free interval, are even more effective in reducing the symptoms of primary dysmenorrhea. In a multicentric observational trial, 406 women suffering from primary dysmenorrhea were treated with continuously administered desogestrel 75 µg/day (Cerazette®) over a period of three cycles. This treatment reduced the frequency of moderately severe to severe dysmenorrhea from 84 % to 7 % (Fig. 2).

In 51 % of women, the symptoms were completely abolished and 39 % observed a light improvement [8]. These improvements are partly due to ovulation inhibition and continuous suppression of the gonadal axis.

### ■ Premenstrual syndrome (PMS)


#### Frequency and symptoms

In the gynaecological consultation, the treatment of women with premenstrual syndrome plays a significant role. The women affected are usually between 30 and menopausal age. Premenstrual syndrome occurs in up to 80 % of women. Every fourth woman has strong complaints in the sense of PMS. In approximately 5 % of women, the symptoms are so strong that the disease is termed premenstrual dysphoric disorder, PMDD [17–20]. It refers to somatic as well as vegetative complaints symptoms (Tab. 2). The symptoms are diverse and the expression varies greatly.

In some of these patients, the complaints are so prominent that they are of clinical importance. Women are often unable to work and also not capable of fulfilling their daily task in the household and in the family.

It is likely that there is a certain disposition for this reaction to the normal fluctuations in the biphasic cycle. The physiological hormonal fluctuations





**Figure 2:** Severe dysmenorrhea before and after 3 month therapy with an estrogen-free pill (0.75 mg desogestrel; Cerazette®), n = 406

after ovulation act as a trigger and lead to the disturbance of the serotonergic system. Serotonin and its various receptors in the CNS are the essential modulators of mood and emotional state. Disturbance leads to a reduced effect of serotonin and to a decreased sensitivity of the GABA-benzodiazepin receptor and the excretion of  $\beta$ -Endorphin.

Therapeutic approaches are the inhibition of ovulation and thus, the cycle-dependent fluctuations. This is possible with any ovulation inhibitor, however, best when used in an extended-cycle or continuous cycle intake. This has been proven by many observational studies with combination preparations [7, 16, 21].

Although the effect of estrogen-free preparations for the treatment of PMS in clinical observational studies has been proven, the only oral preparation currently available for this indication is the combination preparation YAZ® (ethinyl estradiol 20  $\mu$ g and drospirenone 3 mg). It has been approved in the USA, and is taken for 24 days followed by a four-day pill-free interval [22]. However, also the ovulation inhibitory progestin only systems, which all have continuous serum hormone levels, have proven successful in the prevention of PMS and PMDD.

In a placebo controlled crossover study, 43 healthy women with PMS were treated with medroxyprogesterone acetate. A significant improvement of the

symptom scores ( $p < 0.05$ ) was achieved compared to the placebo [23].

This was confirmed by the data of a multicentric, prospective, double-blind, randomized, parallel group study. Here, the patients received progesterone vaginally or rectally [24]. Similar results were observed by Wyatt et al. [25]. In this study, a significant clinical improvement of PMS compared to placebo occurred with the treatment with progesterone (Odds Ratio [OR] = 1.05; 95 %-CI: 1.03–1.08) and also the synthetic progestins (OR = 1.07; 95 %-CI: 1.03–1.11).

In a prospective, six months observational study with women suffering from PMS, Ahrendt compared the effectiveness of the progestin-only pill containing desogestrel 75  $\mu$ g/day (Cerazette®) (n = 33) with the long-cycle intake of a combination preparation containing desogestrel 150  $\mu$ g/day plus ethinyl estradiol 30  $\mu$ g/day (Marvelon®) [26]. During the six months of the study, the premenstrual syndrome improved significantly with the progestin-only medication ( $p < 0.001$ ). This result was not achieved by the combination preparation ( $p = 0.054$ ). In average, the score results in the Cerazette®-group were reduced from 1.45 points to 0.87 points, while they fell from an average of 1.00 points to 0.73 points in the Marvelon® group. The significant reduction occurred within the sixth months of treatment (Screening at treatment month 3:  $p = 0.531$ ; screening at month 6:  $p = 0.002$ ;

**Table 2:** Symptoms of premenstrual dysphoric disorders (PMDD)

#### Psychological symptoms

- Depressed mood
- Pronounced tension, anxiety
- Pronounced mood swings
- Pronounced irritability
- Subjective poor concentration
- Lethargy/fatigue
- Sleeping disorders
- Feeling overwhelmed or out of control

#### Somatic symptoms

- Mastodynia
- Bloating
- Abdominal pain
- Edemas
- Weight gain
- Headache

treatment month 3 and treatment month 6:  $p = 0.005$ ) [26].

## Menstrual Migraine




Up to 60 % of all women report a time-dependent correlation between the onset of migraine and menstruation [27, 28].

### Definition

- Migraine without aura
- 2 days before menstruation
- Up to 3 days after the onset of menstruation
- Pattern confirmed in 66 % of all cycles.

Menstrual migraine has been included by the International Headache Society (IHS) as an independent diagnosis in the headache and facial pain classification (Headache Classification Subcommittee of the International Headache Society [IHS] 2004) [29].

According to the definition, menstrual migraine is a paroxysmally occurring, single-sided, pulsating headache without aura, which exclusively occurs in between two days before until three days after the onset of menstrual bleeding in at least 66 % of the cycles. Possible accompanying symptoms may be nausea, vomiting, phonophobia and photophobia. The headaches as well as the accompanying symptoms in menstrual migraine are usually of longer duration and more severe than in classical migraine with aura [27, 29–31]. The etiology seems to be the physiological reduction



in serum estrogen and serum progestin levels, which act as a trigger. Initially high estrogen levels are probably a prerequisite for the development of migraine attacks [8, 15].

In a study, Ahrendt et al. [8] compared 30 patients with menstrual migraine, who were treated with the estrogen-free pill containing 75 µg Desogestrel (Cerazette®), with 33 patients that received the combination preparation containing 150 µg desogestrel und 30 µg ethinyl estradiol (Marvelon®) in the extended-cycle intake for six months.

The group studied the reduction of frequency, duration and intensity of menstrual migraine and its accompanying symptoms.

Both medications reduced all three examined parameters (Figs. 3–5) of menstrual migraine. The frequency of headaches was significantly reduced in both medication groups ( $p < 0.001$  each). In both groups, the onset of headaches was reduced by an average of four to 2.5 days per cycle. Notably, the intense pain maxima occurred less frequently. This effect was more pronounced under the

use of Cerazette® and occurred earlier. 25 % of the patients were completely painless with the intake of Cerazette®. In 70 % of patients, the migraine attacks were reduced by more than 50 %.

Also the reduction in headache intensity was more pronounced with the use of Cerazette® compared with Marvelon®. In 67 % of the Cerazette®-patients, pain intensity was reduced by more than 50 %, however, only in 48 % of the patients treated with Marvelon®.

The average headache duration was reduced to under a third in patients treated with Cerazette®, and to 60 % in patients treated with Marvelon®. The results were significant in both groups. The average headache intensity was reduced by over 60 % with Cerazette® and to over 30 % with Marvelon®.

Likewise, the accompanying symptoms of migraine were reduced in patients treated with the combination preparation as well as in those patients with extended-cycle treatment, but particularly in patients with estrogen-free pills. Here, the incidence of nausea was reduced by 77 %, of vomiting by 70 %, of photophobia by 78 % and of phonophobia by 64 % [29].

## Risks and Contraindications in Relation to Combined Oral Contraceptives

In gynaecological practice, we see many patients with diseases and risk factors that represent contraindications for COCPs. Such contraindications are:

- Absolute contraindications
  - Acute or severe liver disease
  - Previous or existent benign or malignant liver tumors
  - Acute diseases of the gall bladder
  - Acute diseases of the pancreas
  - Estrogen-dependent tumors: breast cancer, endometrial cancer
  - Previous or existent thromboembolism
  - Family history of thrombosis
  - Thrombophilia
  - Lupus erythematosus, hyperhomocysteinemia, antiphospholipid antibodies
  - Myocardial infarction and stroke
  - Marfan syndrome
  - Severe diabetes mellitus with arteriosclerosis

**Table 3:** Prevalence of hereditary thrombophilia

Risk factors	RR without OC	RR with OC	Prevalence (%)
Clinically healthy women	1	4	0.01–0.02
Pregnancy	5		0.05
Postpartal phase	25		0.25
Family history	3	11	
APC-resistance	8	35	3–5
Protein-C-deficiency	9	15	0.1–0.5
Protein-S-deficiency	8	8	0.1–0.5
Antithrombin-III-deficiency	4	32	0.02–0.05
Antiphospholipid-antibodies			2–9

APC: activated Protein C; OC: oral contraceptives; RR: relative risk

- Idiopathic hyperlipidemia
- Age > 40 years + hypertension + obesity + smoking
- Relative contraindications
  - State after hepatitis
  - Cholestatic liver dysfunction
  - Porphyria
  - Thrombophlebitis
  - Pronounced varicosis
  - Hypertension
  - Otosclerosis

The risk for venous thromboembolisms is particularly increased when taking estrogen-progestin combination preparations [32, 33] (Tab. 3).

If a patient has any of the above-mentioned risk factors or diseases, progestin only systems are usually applicable. In any case, a sensible individual risk-benefit assessment has to be carried together whereby thoroughly informing the patient, which has to be documented.

## ■ Estrogen-free Pills

Of all estrogen-free preparations, the oral preparations (progestin only pills, POPs) are most frequently utilized, as they represent a non-invasive and reversible method of contraception [3]. We differentiate between 3 groups of estrogen-free pills:

- Classical minipill without ovulation inhibition
- Estrogen-free pill with ovulation inhibition
- Postcoital pill for emergency contraception

### Mini pills

Mini pills are pills containing 0.03 mg levonorgestrel that do not inhibit ovula-

tion. This includes the preparations 28 mini®, Microlut® and Micro-30 Wyeth®.

The mechanism of action affects the 3 peripheral parameters: the fallopian tubes (disturbed transport of the ovum), the endometrium (inhibition of nidation) and the cervix (thickening of cervical mucus). In order to achieve high contraceptive efficacy, the minipill has to be taken to a strict time every day. This fact, in turn, reduces the Pearl Index. The safety window is 3 hours. The minipill is usually indicated during lactation.

### Estrogen-free Pills

The so-called estrogen-free pills are preparations with 0.075 mg Desogestrel (Cerazette®). This pill achieves complete ovulation inhibition and, in addition, has an effect via the three peripheral parameters fallopian tube, endometrium and cervix. Contraceptive efficacy is very good and comparable to that of combination preparations. Also the intake scheme is similar to the combination preparations. The safety margin is 12 hours.

Thus, estrogen-free pills are suitable for women of all ages that desire quickly reversible oral contraceptives. In addition, they are suitable for patients with specific indications such as:

- Cycle-dependent complaints: menstrual migraine, premenstrual syndrome (PMS), pelviphathia spastica, dysmenorrhea, hypermenorrhea
- Estrogen-dependent complaints: obesity, tendency for edemas, uterine fibroids, and endometriosis
- Lactation
- Risk factors or diseases, which represent contraindications for estrogen-progestin combination preparations

The estrogen-free pill is well tolerated. Due to the partial androgenic activity of desogestrel, the pill is not suitable for women with seborrhoea, blemished skin or acne. The experience from clinical practice has shown that they are also not suitable for women with recurrent menstrual disorders [34].

Similar to other progestin preparations with continuous intake, irregular bleedings are not predictable. Some patients experience regular weak bleedings, while others suffer from intermediate bleedings, breakthrough bleedings or amenorrhea.

Even when taking the estrogen-free pill Cerazette® for several years, the endogenous estradiol production is not greatly suppressed to an extent that would favour osteoporosis.

### Postcoital pills

For postcoital contraception (Fig. 6), there are two preparations with 750 µg levonorgestrel each (Unofem® and Levogynon®) and one with ulipristal (EllaOne®).

Unofem® and Levogynon® (750 µg levonorgestrel each) are taken once within the first 72 hours *post coitum*. Effectiveness decreases linearly from 12 hours onwards. In total, pregnancy rate is reduced by 85 % (74–93 %) with 750 µg levonorgestrel [35].

Depending on the time point of intake within the cycle, ovulation inhibition or delay can be suspected via the impact on the hypothalamic-pituitary-ovarian axis.

EllaOne® contains Ulipristal, a progesterone receptor modulator of the second generation. Ulipristal acetate is the first approved selective progesterone receptor modulator (SPRM) for emergency contraception. Its exclusive indication is emergency contraception for up to 120 hours (5 days) after unprotected sexual intercourse or in cases of contraceptive failure.

The main mechanism of action is the inhibition or delay of ovulation, but also the effects on the endometrium. The results of two studies show that ulipristal acetate is at least as effective as levonorgestrel in emergency contraception. Both studies showed a significant im-




Figure 6: Postcoital contraception – intake post-intercourse (LNG: levonorgestrel)

provement of the contraceptive effect compared to levonorgestrel, independent of when during the five days the pill was taken. Within the first 0–72 hours: 85 % (Ulipristal) versus 69 % (750 µg levonorgestrel twice); on day 3: 93 % versus 50 % [4]. This preparation was introduced to the market in October 2009.

## ■ Depot injections

Depot injections consist of progestins, which are injected i. m. and are then released protractedly.

- 150 mg medroxyprogesterone acetate (DepoClinovir®): injected every three months (plus/minus 14 days) i.m., starting on the first day of the cycle. The microcrystalline suspension forms a depot, from which continuous resorption occurs. Within the first weeks, the plasma levels range from 1000 to 6000 pg/ml and fall to 500 pg/ml in the last week.
- 200 mg norethisterone enanthate (Noristerat®): initially injected every two months, thereafter every three months i.m. It is stored in the fatty tissue. At the beginning, serum hormone levels are relatively high, and then drop continuously.

## Effect

Depot progestins have a high contraceptive efficacy (Pearl Index 0.1–0.6). The effect is due to ovulation inhibition as well as to the effects on the cervical mucus, the endometrium and the tube motility.

## Cycle

The cycle shows a very variable, unforeseeable bleeding pattern with the administration of depot progestins. Amenorrhea can exist from the beginning onwards, but also irregular or breakthrough bleedings of different severity can occur. The bleeding pattern greatly determines compliance.

If the patient still wants to conceive a child, she has to be informed that the re-establishment of a normal ovulatory cycle can take several months, in some cases up to a year.

## Side effects

Side effects are acne, mastodynia (indication for POPs, see above), headaches and frequently pronounced weight gain.

However, in clinical practice, the long-term intake of MPA can cause bone density loss, as the bone metabolism correlates with the serum estrogen levels. This is especially important in adolescents and young adults, as this is the predominant phase of bone growth. Bone density loss increases with intake duration. It is not known whether MPA reduces bone density in growing women (aged 12–18 years) and whether osteoporosis associated fractures are thus increased later in life [36, 37].

## Advantages

Depot progestins are especially indicated in patients, who desire long-term contraception and who do not want to take a daily

tablet. They are also indicated in patients with contraindications for estrogens, in sickle cell anaemia and during lactation.

Positive effects are also expected in patients with endometriosis, fibroid uterus, dysmenorrhea and hypermenorrhea.

## ■ Implants

As an implant, Implanon® is available. Implanon® is a little rod of 4 cm length and 2 mm diameter. It consists of the medical synthetic material Evatane® and the core contains the progestin etonogestrel (3-ketodesogestrel), the active metabolite of desogestrel. Implanon® is applied subdermally on the first day of the cycle on the upper arm and remains there for three years. Afterwards, it is removed by incision.

## Pharmacokinetics

Within a few days, the serum hormone levels reach their maximum and then level out above the ovulation inhibition dose of 90 pg/ml. Within the first two years, 35–45 µg etonogestrel per day are released; in the 3<sup>rd</sup> year 25–30 µg per day [38].

Implanon® is a very safe method. With a Pearl Index of 0.14, it is comparable to the available oral contraceptives. Apart from ovulation inhibition, also the cervix, the endometrium and the fallopian tubes are affected.

After removal of Implanon®, the woman quickly regains fertility. Already after four days, her serum hormone levels reach normal levels.

The endogenous estradiol levels lie in the prefollicular area. Studies have shown that the bone density is not reduced with Implanon®.

## Menstrual cycle

The bleeding pattern varies greatly with Implanon®. In some patients, a longer amenorrheic phase occurs immediately after implantation. In others, weak bleedings may occur regularly or with greater intervals.

Approximately 15 % have additional or continuous menstrual bleeding to varying degrees, which may lower compliance. However, most irregular bleedings can be treated.



In a study by Mansour et al. [39], the use of Implanon® resulted in the following bleeding pattern: amenorrhea (22.2 %), infrequent bleeding (33.6 %), frequent bleeding (6.7 %) and/or prolonged bleeding (17.7 %) [39, 40].

### Acceptance

The implantation as well as the subsequent removal of the implant is usually uncomplicated. The comfort for the patient is good. Migration of the implants are improbable when correctly placed. In order to correctly position the implant, application has to be learned. Once in a while, implants have been difficult to retrieve due to a too deep positioning.

In the foreseeable future, an x-ray positive implant will be introduced to the market, which will facilitate the retrieval of impalpable implants. Furthermore, a new applicator will be available, which prevents deep insertion of the implant.

If applicable, the tolerance and acceptance of the continuous treatment with desogestrel should be tested with pretreatment with Cerazette® for three months. Also, competitive athletes can use Implanon®.

### Indications

Implanon® is suitable for patients who wish for a long-term contraception, in order not to have to take a daily pill. It is also indicated for patients with cycle-dependent disorders (menstrual migraine, PMS, dysmenorrhea) or estrogen-related disorders (mastodynia, tendency for edemas, endometriosis) and for patients, who lack compliance or who often transgress the time zones.

Implanon® is also approved for contraception during lactation [41].

### Contraindications

Clinical practice has proven that Implanon® is not suitable for patients with acne, hypermenorrhea and recurrent breakthrough bleedings.

## ■ Intrauterine Hormonal Contraception

As an intrauterine hormonal contraceptive, the progestin-containing intrauterine device Mirena® is available, which contains levonorgestrel. Mirena® can remain in the uterus for as long as five

years. Contraceptive efficacy is very high (Pearl Index: 0.14)

### Pharmacokinetics

Not long after the insertion of Mirena®, constant serum hormone levels of 100–200 pg/ml are reached.

Contraceptive efficacy of Mirena® relies on the local effects of levonorgestrel: Thickening of the cervical mucus, endometrial alterations and fallopian tube effects. Ovulation is not inhibited, as the ovulation inhibition dose in intrauterine release of levonorgestrel is 50 µg/day but Mirena® only releases 20 µg/day.

### Menstrual cycle

The bleeding pattern after Mirena® implantation varies greatly. Some patients experience the immediate onset of amenorrhea, while others have regular and weak bleedings. In approximately 30 % of women, irregular and breakthrough bleeding occur within the first three to six months, which are of variable duration and intensity. However, they can usually be treated by the additional administration of levonorgestrel-containing pills for two months or one tablet of ethinyl estradiol for 21 days) [42]. In the majority of patients, however, bleeding intensity is greatly reduced with a positive influence on hemoglobin and ferritin and thus on the wellbeing and the performance of the patient [43, 44].

### Advantages

Contraceptive efficacy remains for a period of five years. Mirena® is compliance-independent and acceptance is high [45]. This is partly explained by the fact that Mirena® causes amenorrhea lasting for several months in many patients. This, in turn, diminishes cycle-dependent disorders and the non-contraceptive additional effect concerning these cycle dependent complaints (menstrual migraine, abdominal pain, PMS but also mastodynia) is very high. Mirena® is especially indicated in patients with the wish for long term contraception, in cycle-dependent disorders, especially dysmenorrhea, hypermenorrhea and perimenopausal bleeding disorders, which has been confirmed by various studies [42, 46]. Due to the local progestin effect, estrogen monotherapy can be administered in menopausal complaints, which reduces the systemic effect of progestin components compared to the combina-

tion preparations. Mirena® is also indicated during lactation.

### Side effects

Breakthrough bleedings can occur within the first three months of application, thereafter, they occur rather seldom. Clinical experience shows that disturbing side effects, which are often a reason for treatment termination, are relatively rare. Amenorrhea is often seen as an advantage and is often wished for. The frequency of mastodynia, weight gain and skin alterations confer to the general population.

The risk for ascending infections is reduced compared to the general population. However, the risk is significantly lower compared to the copper IUD [47, 48].

### Treatment of Irregular Breakthrough Bleedings with Estrogen-free hormonal Contraceptives

It is a disadvantage that due to a continuous use without intervals in all preparations, cycle control remains difficult. Periodic bleedings or weak bleedings may occur but also oligomenorrhea and amenorrhea. However, very often, breakthrough bleedings of varying degree, duration and frequency occur, which greatly affect the patients and sometimes even are a reason for the discontinuation of the treatment. However, these side effects are reduced with longer intake of the preparations.

For the everyday practice, it is important to know the individual characteristics of the irregular bleedings and, especially, their treatment.

### Therapeutic management of bleeding disorders

As evidence based studies are not yet available on this subject, experience and opinions of expert panels are of great importance. Primarily, the following questions should be answered (according to [49]):

- Do the irregular bleedings occur within the first three months or later?
- Is the endometrium high (> 6 mm), low or maybe atrophic?

Within the first 3 months, hormonal dysregulation is to be expected. The continuous estrogen-free medication causes



**Table 4:** Therapy of irregular bleedings within the first three months (according to [49])

Endometrial thickness	Options	Therapy
Thin ( $\leq 6$ mm)	Option 1	One tablet ethinyl estradiol 25 µg for 21 days or 50 µg E2-patch for 4–7 days
	Option 2	Tranexam acid (Cyklokapron®): 3 × 1 tablet for 5 days
Thick ( $> 6$ mm)	Option 1	1 tablet norethisterone acetate 2.5–5 mg for 10 days
	Option 2	Estrogen-free pills: discontinuation of pill for 4 days

**Table 5:** Therapy of irregular bleedings after more than three months of intake (according to [49]).

Endometrial thickness	Options	Therapy
Thin ( $\leq 6$ mm)	Option 1	Doxycyclin 100 mg for 10 days
	Option 2	Tranexam acid (Cyklokapron®): 3 × 1 tablet for 5 days
	Option 3	Ethinyl estradiol 25 µg for 21 days or 50 µg E2-patch for 4–7 days
Thick ( $> 6$ mm)	Option 1	1 tablet norethisterone acetate 2.5–5 mg for 10 days
	Option 2	With estrogen-free pill and an implant: additional tablet Cerazette® for 28 days
	Option 3	Estrogen free pills: discontinuation of pill for 4–7 days

ovary suppression as well as a lack of proliferation and irregular secretion [40]. Here, hormonal therapy is the first choice, depending on endometrial thickness.

#### Irregular bleedings within the first three months

In irregular bleedings within the first 3 months, Ahrendt et al. [49] suggest the following therapeutic options outlined in Table 4.

#### Irregular bleedings after more than three months of intake

Later occurring bleedings are probably due to the vascular compartment of the endometrium, i. e. angiogenetic disorders in atrophic endometrium [50]. Mechanisms that have been discussed are the suppressed development of the endometrial spiral arteries by progesterone and the subsequent altered homeostasis [51].

Increased vein development and dilation at the endometrial transition zone [40, 52, 53] as well as an alteration of the leucocytes in the sense of an inflammatory process [54, 55]. This increasingly causes microvascular fragility and thus irregular bleedings. In these cases of mostly atrophic endometrium, treatment with tranexem acid [40] or the prosta-

acid [56] has led to longer bleeding free periods in studies as well as in clinical practice. Both substances can increase vascular stability. As subclinical endometritis [55] is also discussed as a cause for the bleedings, treatment with doxycyclin might be effective.

As an exact cause for irregular bleeding with the use of estrogen-free systems can usually not be determined, the individual therapy has to be based on hypotheses and knowledge from single studies and clinical experience.

In irregular bleedings after an intake of more than three months, the following therapies, listed in Table 5, are possible [49].

With a strict adherence to these therapy options, disturbing breakthrough bleedings with the use of progestin-only systems can be reduced to a minimum. This then prevents therapy discontinuation and significantly improves compliance.

### ■ Conflict of interest

The author declares that there is no conflict of interest as defined by the guidelines of the International Committee of Medical Journal Editors (ICMJE; www.icmje.org).

### References:

- Jick H, Dinan B, Herman R, Rothman KJ. Myocardial infarction and other vascular diseases in young women. Role of estrogens and other factors. *JAMA* 1978; 240: 2548–52.
- Thorogood M, Vessey MP. An epidemiologic survey of cardiovascular disease in women taking oral contraceptives. *Am J Obstet Gynecol* 1990; 163: 274–81.
- Ahrendt HJ. Hormonale Kontrazeption. *CME Prakt Fortbild Gynakol Geburtsmed Gynakol* 2009; 5: 184–93.
- Rabe T, Ahrendt HJ, König K, Ludwig M, Goeckenjan M, Merkle E, Thonke I, Zahradnick HP. Postkoitale Kontrazeption. *Frauenarzt* 2009; 50: 784–90.
- Hatcher RA, Guillebaud J. The pill: combined oral contraceptives. In Hatcher RA et al. (eds). *Contraceptive Technology*. 17th ed. Ardent Media, Inc, New York, 1998; 405–60.
- Oddens BJ, Arnolds HT, Van Maris MG, Van Lunsen HW. The dynamics of oral contraceptive use in The Netherlands 1990–1993. *Adv Contracept* 1994; 10: 167–74.
- Kuhl H, Jung-Hoffmann C. Kontrazeption. Thieme, Stuttgart, New York; 1999.
- Ahrendt HJ, Kleinschmidt S, Kropf S, Kose D. Präventive Wirkung hormoneller Kontrazeptiva im Langzyklus bei menstrueller Migräne. *Frauenarzt* 2007; 48: 1186–92.
- Ahrendt HJ, Apetauer I. Improvement of oestrogen-related symptoms with a desogestrel-only contraceptive among women who experienced oestrogen-related symptoms using combined oral contraceptives [Abstract P100]. *Eur J Contracept Reprod Health Care*; 2008; 13: 102–3.
- Dawood MY. Primäre Dysmenorrhö: Fortschritte in der Pathogenese und Management. *Geburtsh Gynakol* 2006; 108: 428–41.
- Bieglmayer C, Hofer G, Kainz C, Reinthaller A, Kopp B, Janisch H. Concentrations of various arachidonic acid metabolites in menstrual fluid are associated with menstrual pain and are influenced by hormonal contraceptives. *Gynecol Endocrinol* 1995; 9: 307–12.
- Proctor ML, Roberts H, Farquhar CM. Kombinierte Pille (OCP) für die Behandlung primärer Dysmenorrhoe. *Cochrane Database Syst Rev*. 2001; (4): CD002120.
- Rosenwaks Z, Seegar-Jones G. Menstrual pain. Its origin and pathogenesis. *J Reprod Med* 1980; 25: 207–12.
- Hendrix SL, Alexander NJ. Primäre Dysmenorrhoe Behandlung mit Desogestrel-haltige niedrig dosierte orale Kontrazeptiva. *Empfängnisverhütung* 2002; 66: 393–9.
- Edelman AB, Gallo MF, Jensen JT, Nichols MD, Schulz KF, Grimes DA. Continuous or extended cycle vs cyclic use of combined oral contraceptives for contraception. *Cochrane Database Syst Rev*. 2005(3): CD004695.
- Göretzlehner G. Langzyklus – Langzeiteinnahme, Anwendung des Dienogestrel-haltigen oralen hormonalen Kontrazeptivums, der Mikropille ValetteR. Muhlheim/Ruhr: H.U.F 2009.
- Angst J, Sellaro R, Merikangas KR, Endikott J. The epidemiology of perimenstrual psychological symptoms. *Acta Psychiatr Scand* 2001; 104: 110–16.
- Wittchen HU, Becker E, Lieb R, Krause P. Prevalence, incidence and stability of premenstrual dysphoric disorder in the community. *Psychol Med* 2002; 32: 119–32.
- Cohen LS, Miner C, Brown EW, Freeman E, Halbreich U, Sundell K, McCray S. Premenstrual daily fluoxetine for premenstrual dysphoric disorder: a placebo controlled, clinical trial using computerized diaries. *Obstet Gynecol* 2002; 100: 435–44.
- Hylan TR, Sundell K, Judge R. The impact of premenstrual symptomatology on functioning and treatment-seeking behavior: experience from the United States, United Kingdom and France. *J Womens Health Gend Based Med* 1999; 8: 1043–52.
- Bachmann G, Sulak PJ, Sampson-Landers C, Benda N, Marr J. Efficacy and safety of a low-dose 24-day combined oral contraceptive containing 20 micrograms ethinylestradiol and 3 mg drospirenone. *Contraception* 2004; 70: 191–8.
- Pearlstein TB, Bachmann GA, Zacur HA, Yonkers KA. Treatment of premenstrual dysphoric disorder with a new drospirenone-containing oral contraceptive formulation. *Contraception* 2005; 72: 414–21.
- Hellberg D, Claesson B, Nilsson S. Premenstrual tension: a placebo-controlled efficacy study with spironolactone and medroxyprogesterone acetate. *Int J Gynaecol Obstet* 1991; 34: 243–8.
- Magill PJ. Investigation of the efficacy of progesterone pessaries in the relief of symptoms of premenstrual syndrome. *Progesterone Study Group. Br J Gen Pract* 1995; 45: 589–93.

25. Wyatt K, Dimmock P, Jones P, Ohrai M, O'Brien S. Efficacy of progesterone and progestogens in management of premenstrual syndrome: systematic review. *BMJ* 2001; 323: 776–80.
26. Ahrendt HJ. Advantages and challenges of estrogen-free hormonal contraception. *BJOG* 2009 (in press).
27. Couturier EG, Bomhof MA, Neven AK, van Duijn NP. Menstrual migraine in a representative Dutch population sample: prevalence, disability and treatment. *Cephalalgia* 2003; 23: 302–8.
28. Allais G, Benedetto C. Updates on menstrual migraine: from clinical aspects to therapeutic strategies. *Neurol Sci* 2004; 25 (Suppl 3): 229–31.
29. Headache Classification Subcommittee der International Headache Society (IHS). Die Internationale Klassifikation von Kopfschmerzkrankungen. 2. Aufl. *Cephalalgia* 2004; 24 (Suppl 1): 1–160.
30. Granello F, Sances G, Allais G, Nappi RE, Tirelli A, Benedetto C, Brundu B, Facchinetti F, Nappi G. Characteristics of menstrual attacks in women with menstrually related migraine referred to headache centres. *Cephalalgia* 2004; 24: 707–16.
31. MacGregor EA. "Menstrual" migraine: towards a definition. *Cephalalgia* 1996; 16: 11–21.
32. Bauersachs R, Lindhoff-Last E, Ehrly A M, Kuhl H. Die Bedeutung der hereditären Thrombophilie für das Thrombose-risiko unter der oralen Kontrazeption. *Zbl Gynakol* 1996; 118: 262–70.
33. Koster T, Small RA, Rosendaal FR, Helmerhorst FM. Oral contraceptives and venous thromboembolism: a quantitative discussion of the uncertainties. *J Intern Med* 1995; 238: 31–7.
34. Ahrendt HJ, Kose D. Östrogenfreie Pille: Blutungsverhalten und Therapie von Zusatzblutungen. *Frauenarzt* 2008; 49: 134–40.
35. Task Force on Postovulatory Methods of Fertility Regulation. Randomised controlled trial of levonorgestrel versus the Yuzpe regimen of combined oral contraception for emergency contraception. *Lancet* 1998; 352: 428–33.
36. Noristerat Fachinformation. BPI Service GmbH FachInfo-Service, Aulendorf.
37. Depoclinovir Fachinformation. BPI Service GmbH. FachInfo-Service, Aulendorf.
38. Olsson SE, Odland V, Johansson E. Clinical results with subcutaneous implants containing 3-keto desogestrel. *Contraception* 1990; 42: 1–11.
39. Mansour D, Korver T, Marintcheva-Petrova M, Fraser IS. The effects of ImplanonR on menstrual bleeding patterns. *Eur J Contracept Reprod Health Care* 2008; 13: 13–28.
40. Hickey M, d'Arcangues C. Vaginal bleeding disturbances and implantable contraceptives. *Contraception* 2002; 65: 75–84.
41. Reinprayoon D, Taneepanichskul S, Bunyavejchevin S, Thaihumyanon P, Punnahitananda S, Tosukhowong P, Machielsen C, van Beek A. Effects of the etonogestrel releasing contraceptive implant (ImplanonR) on parameters of breastfeeding compared to those of an intrauterine device. *Contraception* 2000; 62: 239–46.
42. Busfield R, Farquhar C, Sowter M, Lethaby A, Sprecher M, Yu Y, Sadler L, Brown P, Johnson N. A randomised trial comparing the levonorgestrel intrauterine system and thermal balloon ablation for heavy menstrual bleeding. *BJOG* 2006; 113: 257–63.
43. Allonen H, Kulmala Y. Return to fertility after the removal of Nova-T or the Levonorgestrel-IUD. *Leiras study report* 1991; 1205: 1–23.
44. Andersson K, Rybo G. Levonorgestrel-releasing intrauterine device in the treatment of menorrhagia. *Gynaecol* 1990; 97: 690–4.
45. Mirena® Satisfaction Study. Global Market Research. Bayer HealthCare. May 2008.
46. de Jonge ET, Yigit R, Molenberghs G, Straetmans D, Ombelet W. Predictors of oligoamenorrhea at 1-year follow up in premenopausal women using a levonorgestrel-releasing intrauterine system. *Contraception* 2007; 76: 91–5.
47. Andersson K, Odland V, Rybo G. Levonorgestrel-releasing and copper-releasing IUDs during five years of use: A randomized comparative trial. *Contraception* 1994; 49: 56–72.
48. Römer T. Erfahrungen und Empfehlungen zur Anwendung des Levonorgestrel-Intrauterinsystems. *Thieme Praxis Report* 2009; 1: 1–24.
49. Ahrendt et al. 2008. Bibliographische Angaben bitte ergänzen.
50. Römer T. Blutungsstörungen unter Ovulationshemmern. *Gynecol Endocrinol* 2007; 5: 66–70.
51. Johannisson E. Endometrial morphology during the normal cycle and under the influence of contraceptive steroids. In: d'Arcangues C, Fraser IS, Newton JR, Odland V (eds). *Contraception and Mechanisms of Endometrial Bleeding*. Cambridge: WHO. Cambridge University Press 1990; 53–81.
52. Kovacs G. Progestogen-only pills and bleeding disturbances. *Hum Reprod* 1996; 11: 20–3.
53. Taubert HD, Kuhl H (Hrsg). *Kontrazeption mit Hormonen*. 2. Aufl. Thieme, Stuttgart, 1995.
54. Song JY, Russell P, Markham R, Manconi F, Fraser IS. Effects of high-dose progestagens on white cells and necrosis in human endometrium. *Human Reprod* 1996; 11: 1713–8.
55. Vincent AJ, Salomonson LA. MMP leucocytes and steroid associated uterine bleeding. *Human Reprod* 2000; 15: 135–44.
56. Kaewrudee S, Taneepanichskul S, Jaisamraun U, Reinprayoon D. The effect of mefenamin acid on controlling irregular uterine bleeding secondary to Norplant use. *Contraception* 1999; 60: 25–30.

### Further References:

- Ahrendt HJ, Karck U, Pichl T, Müller T, Ernst U. The effects of estrogen-free, desogestrel-containing oral contraceptive in women with cyclical symptoms: Results from two studies on oestrogen-related symptoms and dysmenorrhoea. *Eur J Contracept Reprod Health Care* 2007; 12: 354–61.
- Pomp ER, le Cessie S, Rosendaal FR, Doggen CJ. Risk of venous thrombosis: obesity and its joint effect with oral contraceptive use and prothrombotic mutations. *Brit J Haematol* 2007; 139: 289–96.
- Visser WH, Jaspers NM, de Vriend RH, Ferrari MD. Risk factors for headache recurrence after sumatriptan. *Cephalalgia* 1996; 16: 264–9.

# Mitteilungen aus der Redaktion

Besuchen Sie unsere Rubrik

## ☒ Medizintechnik-Produkte



Neues CRT-D Implantat  
Intica 7 HF-T QP von Biotronik



Artis pheno  
Siemens Healthcare Diagnostics GmbH



Philips Azurion:  
Innovative Bildgebungslösung

Aspirator 3  
Labotect GmbH



InControl 1050  
Labotect GmbH

## e-Journal-Abo

Beziehen Sie die elektronischen Ausgaben dieser Zeitschrift hier.

Die Lieferung umfasst 4–5 Ausgaben pro Jahr zzgl. allfälliger Sonderhefte.

Unsere e-Journale stehen als PDF-Datei zur Verfügung und sind auf den meisten der marktüblichen e-Book-Readern, Tablets sowie auf iPad funktionsfähig.

## ☒ Bestellung e-Journal-Abo

### Haftungsausschluss

Die in unseren Webseiten publizierten Informationen richten sich **ausschließlich an geprüfte und autorisierte medizinische Berufsgruppen** und entbinden nicht von der ärztlichen Sorgfaltspflicht sowie von einer ausführlichen Patientenaufklärung über therapeutische Optionen und deren Wirkungen bzw. Nebenwirkungen. Die entsprechenden Angaben werden von den Autoren mit der größten Sorgfalt recherchiert und zusammengestellt. Die angegebenen Dosierungen sind im Einzelfall anhand der Fachinformationen zu überprüfen. Weder die Autoren, noch die tragenden Gesellschaften noch der Verlag übernehmen irgendwelche Haftungsansprüche.

Bitte beachten Sie auch diese Seiten:

**Impressum**

**Disclaimers & Copyright**

**Datenschutzerklärung**