

Journal für

Reproduktionsmedizin und Endokrinologie

– Journal of Reproductive Medicine and Endocrinology –

Andrologie • Embryologie & Biologie • Endokrinologie • Ethik & Recht • Genetik
Gynäkologie • Kontrazeption • Psychosomatik • Reproduktionsmedizin • Urologie



Clinical Applications for Estetrol

Visser M, Coelingh Bennink HJT

J. Reproduktionsmed. Endokrinol 2010; 7 (Sonderheft

1), 56-60

www.kup.at/repromedizin

Online-Datenbank mit Autoren- und Stichwortsuche

Offizielles Organ: AGRBM, BRZ, DVR, DGA, DGGEF, DGRM, DIR, EFA, OEGRM, SRBM/DGE

Indexed in EMBASE/Excerpta Medica/Scopus

Krause & Pachernegg GmbH, Verlag für Medizin und Wirtschaft, A-3003 Gablitz

Datenschutz:

Ihre Daten unterliegen dem Datenschutzgesetz und werden nicht an Dritte weitergegeben. Die Daten werden vom Verlag ausschließlich für den Versand der PDF-Files des Journals für Pneumologie und eventueller weiterer Informationen das Journal betreffend genutzt.

Lieferung:

Die Lieferung umfasst die jeweils aktuelle Ausgabe des Journals für Pneumologie. Sie werden per E-Mail informiert, durch Klick auf den gesendeten Link erhalten Sie die komplette Ausgabe als PDF (Umfang ca. 5–10 MB). Außerhalb dieses Angebots ist keine Lieferung möglich.

Abbestellen:

Das Gratis-Online-Abonnement kann jederzeit per Mausklick wieder abbestellt werden. In jeder Benachrichtigung finden Sie die Information, wie das Abo abbestellt werden kann.

Das e-Journal **Journal für Pneumologie**

- ✓ steht als PDF-Datei (ca. 5–10 MB) stets internetunabhängig zur Verfügung
- ✓ kann bei geringem Platzaufwand gespeichert werden
- ✓ ist jederzeit abrufbar
- ✓ bietet einen direkten, ortsunabhängigen Zugriff
- ✓ ist funktionsfähig auf Tablets, iPads und den meisten marktüblichen e-Book-Readern
- ✓ ist leicht im Volltext durchsuchbar
- ✓ umfasst neben Texten und Bildern auch eingebettete Videosequenzen.

Clinical Applications for Estetrol*#

M. Visser, H. J. T. Coelingh Bennink

In this paper the potential clinical applications for the human fetal estrogen estetrol (E_4) are presented based on recently obtained data in preclinical and clinical studies. In the past E_4 has been classified as a weak estrogen due to its rather low estrogen receptor affinity. However, recent research has demonstrated that due to its favorable pharmacokinetic properties, especially the slow elimination and long half-life, E_4 is an effective orally bioavailable estrogen agonist with estrogen antagonistic effects on the breast in the presence of estradiol. Based on the pharmacokinetic properties, the pharmacological profile and the safety and efficacy results in human studies, E_4 seems potentially suitable as a drug for human use in applications such as hormone replacement therapy (vaginal atrophy and vasomotor symptoms), contraception, osteoporosis and breast cancer **J Reproduktionsmed Endokrinol 2010; 7 (Special Issue 1): 56–60.**

Key words: estetrol, E_4 , clinical applications

■ Introduction

Estetrol (E_4) is a human steroid, produced by the fetal liver during pregnancy only. This natural hormone was discovered in urine of pregnant women by Diczfalussy and coworkers in 1965 [1]. Based on its physical and chemical characteristics it was concluded that E_4 is identical with 15α -hydroxyestriol (15α -OHE3) or estra-1, 3, 5(10)-triene-3,15 α -, 16 α -, 17 β -tetrol [2]. Estetrol has the structure of an estrogenic steroid with four hydroxyl groups which explains the acronym E_4 . Estetrol is synthesized by the fetal liver during human pregnancy and reaches the maternal circulation through the placenta. The fetal liver is the exclusive site of 15α - and 16α -hydroxylation [3–5]. After birth the neonatal liver rapidly loses its capacity to synthesize E_4 .

Estetrol was already detected at 9 weeks of pregnancy in maternal urine [6, 7]. During the second trimester of pregnancy high levels were found in maternal plasma, with steadily rising concentrations of unconjugated E_4 to about 1 ng/mL (3 nmol/L) towards the end of pregnancy [8, 9]. So far the physiological function of E_4 has not been studied and is unknown.

The possible use of E_4 as a marker for fetal well-being has been studied quite extensively. However, due to the large intra- and interindividual variation of maternal E_4 plasma levels during pregnancy this appeared not to be feasible

[10–14]. More details on the history of E_4 and data from studies in the period from 1965 to 1984 have been summarized in two review papers [8, 15].

During the last 7 years E_4 has been studied extensively. High oral absorption and bioavailability with a 2–3 h elimination half-life in the rat has been established [16]. In the human E_4 showed a high and dose-proportional oral bioavailability and a remarkably long terminal elimination half-life of about 28 h [17].

Estetrol has a moderate affinity for both human ER α and ER β receptors with a 4–5-fold preference for the ER α [18]. In addition it was found that E_4 binds highly selectively to the estrogen receptors.

In rat and human hepatocytes the rate of E_4 metabolism was studied and found to be slow [18]. In addition E_4 did not inhibit any of the major drug metabolizing cytochrome P450 enzymes CYP1A2, CYP2C9, CYP2C19, CYP2D6 and CYP3A4 at a high concentration of 10 μ M [18]. This in contrast with the effects of estradiol (E_2) and ethinyl-estradiol (EE) on these enzymes.

Using the HepG2 and Hep89 cell lines the ER α -dependent effect of E_2 , estriol (E_3), E_4 and EE on SHBG production was investigated [19]. Estetrol did not stimulate the production of SHBG in both cell lines, while E_2 , E_3 and EE all showed a dose-dependent ER α -mediated increase in SHBG production. Es-

tradiol showed the most prominent increase in the production of SHBG, while E_3 and EE showed a lower and comparable increase in SHBG. In addition no detectable binding of E_4 was found to both the estrogenic and androgenic human SHBG steroid-binding sites [19]. Both testosterone and E_2 were bound with high affinity, whereas EE bound to SHBG with low affinity [19]. These results indicate that the plasma distribution of E_4 or its availability for target tissues may not be affected by SHBG levels in contrast to other natural steroids such as E_2 and testosterone.

Estetrol has also been studied in predictive, validated, pharmacological in vivo rat models and in phases I and IIA clinical trials in the human females. This data indicate that E_4 may be suitable for use in several indications, e. g. hormone replacement therapy (vasomotor symptoms and vaginal atrophy), contraception, osteoporosis and breast cancer. In the current paper these potential clinical applications of E_4 are reviewed based on data generated from pharmacological studies as well as clinical trials performed with E_4 .

■ Vasomotor Symptoms

The efficacy of E_4 in alleviating hot flushes was studied in an experimental rat model considered representative for menopausal vasomotor symptoms [20].

In this model the thermal responses in the tail skin of morphine-dependent

* Reprint of: J Steroid Biochem Mol Biol 2009; 114: 85–9. Reprint with permission from Elsevier. © 2009 Elsevier Ltd. All rights reserved.

Lecture presented at the 18th International Symposium of the Journal of Steroid Biochemistry and Molecular Biology, 18–21 September 2008, Seefeld, Tyrol, Austria.

Correspondence: Monique Visser, Pantarhei Bioscience B. V. Boslaan 11, NL-3701 CH Zeist, The Netherlands; e-mail: mv@pantarheibio.com

ovariectomised (OVX) rats was recorded after administration of naloxone (NAL). Six groups of rats were treated orally for 8 days as follows: vehicle (negative) control; E_4 : 0.1, 0.3, 1.0 and 3.0 mg/(kg day) and as active (positive) control EE: 0.3 mg/(kg day). On day 8, tail skin temperature (TST) was recorded at baseline and for 60 min at 5-min intervals following NAL administration. In control animals TST increased sharply by about 4.5 °C after NAL treatment and reverted to baseline by 60 min. Estetrol suppressed the TST increase in a dose-dependent fashion (Fig. 1).

The highest dose of E_4 tested (3 mg/[kg day]) was equipotent to a 10-fold lower dose of EE. Both fully suppressed TST changes [20].

It is concluded that E_4 is effective in preventing temperature rises in an experimental animal model considered representative for studying the effect of drugs on the menopausal hot flush (vasomotor symptoms). In this model the potency of E_4 was 10-fold lower compared to EE. A phase I multiple dose study to evaluate the safety, tolerability, pharmacokinetics and pharmacodynamics of four dosages of E_4 after daily administration for 28 days was performed in healthy postmenopausal women (unpublished data). In the 2 and 10 mg/day E_4 dose groups women were included with at least 30 hot flushes per week. Estetrol was effective in alleviating hot flushes in most women in both dose groups comparable with a 2 mg/day E_2 -valerate control group during the treatment period of 4 weeks. Although the level of evidence of this study design is not high, it indicates efficacy at these dose levels.

These results suggest that E_4 may be effective for the treatment of hot flushes and other vasomotor symptoms in peri- and postmenopausal women.

■ Vaginal Atrophy

The effect of E_4 on vaginal cornification and uterine weight was studied in OVX rats [21]. Six groups of rats were treated orally once daily for 7 days as follows: vehicle (negative) control; E_4 : 0.1, 0.3, 1.0 and 3.0 mg/(kg day) and EE 0.05 mg/(kg day) as active (positive) control. Vaginal lavages were obtained daily and on day 7 uterine wet weight




Figure 1: The effects of estetrol and ethinylestradiol on the hot flush response induced by naloxone in female ovariectomised rats.




Figure 2: Number of animals with vaginal cornification over a 7-day treatment period with orally administered estetrol (E_4 : 0.1, 0.3, 1.0 or 3.0 mg/(kg day)), ethinylestradiol.

was determined. Vaginal cornification was observed by day 5 in all rats at all E_4 doses and in the animals receiving EE, but not in the control rats (Fig. 2). The onset of cornification with E_4 was dose-dependent. After 7 days treatment, the 2 highest E_4 doses (1.0 and 3.0 mg) induced statistically significantly higher uterine wet weight (myometrium) compared to vehicle [21]. Also a pharmacological study was performed to investigate the effect of E_4 on prevention of bone loss [16]. In this study the uterus of the OVX rats was excised after 4 weeks of treatment. Wet uterine weight (myometrium) was estimated and histological investigation of the endometrium was performed. Four weeks of E_4 treatment

induced dose-dependent increases in uterine weight of OVX rats. The potency of EE in increasing uterine weight in this model was 5–25 times higher than that of E_4 .

In addition, estetrol showed a dose-dependent proliferative estrogenic effect on the rat endometrium after 4 weeks treatment [16]. The order of increasing potency per mg/(kg day) was estimated as follows: 0.1 mg E_4 < 0.5 mg E_4 < 0.1 mg EE < 2.5 mg E_4 , indicating that E_4 was less potent than EE. In summary, estrogenic activity of E_4 was demonstrated in three tissues in OVX rats; vaginal epithelium, myometrium and endometrium. The potency of E_4 was




Figure 3: Estimation of ED₅₀ for ovulation inhibition in 4-day cycling rats treated twice daily with the indicated oral doses of ethinylestradiol (EE) or estetrol (E₄).




Figure 4: The percentage change in plasma levels of luteinising hormone (LH), follicle stimulating hormone (FSH) and sex hormone binding globulin (SHBG) after administration of 2, 10, 20 and 40 mg/day estetrol (E₄) and 2 mg/day estradiol-valerate (E₂) for 28 days.

approximately 20-fold lower compared to EE.

In a phase I multiple dose study with 2, 10, 20 and 40 mg/day E₄ dose groups and a 2 mg/day E₂-valerate control group all administered for 28 days, the effects of E₄ on vaginal cytology and endometrial thickness were investigated (unpublished data). In this study 2 mg/day E₄ for 28 days was as effective as 2 mg/day E₂-valerate in shifting from mainly parabasal cells towards a high percentage intermediate and superficial cells, indicating estrogenic activity of E₄. Proliferation of the endometrium with 2 mg/day E₄ was less than with 2 mg/day E₂-valerate, but 10 mg/day E₄ had a stronger effect.

From these data it can be concluded that E₄ may be suitable for the treatment of urogenital atrophy and the accompany-

ing clinical complaints such as vaginal dryness and dyspareunia in estrogen deficient women. Since E₄ induces proliferation of the endometrium, in women with a uterus, protection against endometrial hyperplasia and cancer may be required, depending on the optimal E₄ dose and the endometrial proliferation induced by that dose.

■ Contraception

The effects of E₄ on ovulation inhibition was studied in regularly cycling female rats [22, 23]. The animals were treated orally twice daily for 4 consecutive days, starting on the day of estrus, with E₄ (0.03, 0.1, 0.3, 1.0 or 3.0 mg/kg), with the comparator EE (0.0003, 0.001, 0.003, 0.01 or 0.03 mg/kg) or with vehicle control. The primary endpoint was the number of ovulated oocytes in the genital tract. Ovulation was inhibited

with E₄ at the twice-daily dose of 0.3 mg/kg and the higher doses (p < 0.05). Ovulation was also significantly inhibited (p < 0.05) by twice-daily administration of the comparator EE at the highest dose (0.03 mg/kg). The ED₅₀ of the dose response curves of EE and E₄ shows that E₄ is about 18 times less potent compared to EE [22] (Fig. 3).

In addition, the effect on plasma levels of gonadotrophins (LH and FSH) was studied in a single rising dose pharmacokinetics study in healthy postmenopausal women with administration of single doses of 0.1, 1, 10 and 100 mg E₄ [17]. LH levels were suppressed in a dose-dependent manner and a profound and sustained inhibition of FSH levels was observed, lasting over 7 days in the 100 mg dose group (FSH was only measured in the 100 mg dose group). In a multiple rising dose study in healthy postmenopausal women with 2, 10, 20 and 40 mg/day E₄ dose groups and a 2 mg/day E₂-valerate control group administered for 28 days, both FSH and LH levels decreased dose-dependently (Fig. 4). From these data it was concluded that E₄ has a profound central inhibitory and dose-dependent effect on gonadotrophins, expected to contribute to the contraceptive effect of E₄. This was further investigated in a phase IIA clinical trial in premenopausal women with proven ovulation in the previous cycle. In this study 4 treatment groups were included: a 10 mg E₄ only group, a 20 mg E₄ only group, a 20 mg E₄/150 µg desogestrel group and a 20 mg E₄/200 mg progesterone group. All treatments were for 28 days. In the E₄/desogestrel group ovulation was inhibited in all women, while in the 10 and 20 mg E₄ only groups in one-third and two-third of the cycles ovulation was inhibited respectively.

With both the 10 and 20 mg E₄ only doses no breakthrough bleeding or spotting occurred, while in the E₄/desogestrel group bleeding was acceptable.

In summary, based on the available data, it can be concluded that E₄ seems suitable to replace EE in combined oral contraceptives.

■ Osteoporosis

In ovariectomised (OVX) rats the bone-sparing effect of oral E₄ was compared to

that of EE [16]. Once-daily oral treatment with a dose of 0.1, 0.5, or 2.5 mg/(kg day) of E_4 or by 0.1 mg/(kg day) of EE as positive control was given for 4 weeks. In this study the following parameters were assessed: (i) serum osteocalcin, (ii) bone mineral density (BMD), bone mineral content (BMC) and bone mineral area (BMA) of lumbar vertebrae L3–L6, (iii) peripheral quantitative computed tomography (pQCT) of the left tibiae and (iiii) the biomechanical properties (strength) of the distal femora. In this rat model E_4 was able to significantly inhibit the OVX-related increases in osteocalcin levels, BMD and BMC, and bone strength (Fig. 5) in a dose-dependent manner. The relative potency of the highest dose of E_4 of 2.5 mg/(kg day) was comparable to the 0.1 mg/(kg day) EE dose, used as positive control. From these data it can be concluded that oral administration of E_4 conveys dose-dependent bone-sparing effects of high quality bone in estrogen depleted OVX rats [16].

In the multiple rising dose study with E_4 in postmenopausal women a dose-dependent decrease of both the marker of bone resorption C-telopeptide and the marker of bone formation osteocalcin was observed (Fig. 6). At the higher dose levels of E_4 (20 and 40 mg) the inhibitory effect on bone formation was only 10 %, whereas bone resorption was suppressed by 35–50 %. This suggests uncoupling of bone metabolism with a preference for bone formation.

These data indicate that E_4 may be suitable as drug for the prevention of osteoporosis in postmenopausal women. It may also be worthwhile to investigate the potency of E_4 to treat osteoporosis and osteoporotic fractures.

■ Breast Cancer

Two prevention studies and one intervention study with E_4 were performed in a rat breast cancer model, in which the animals were treated with DMBA (7,12 dimethylbenz(a)anthracene) to develop estrogen-responsive breast tumors [24]. In the prevention studies the effect on the number and size of the tumors was investigated of oral doses of E_4 in a dose range of 0.5–3.0 mg/kg. The intervention study used oral doses of 1, 3 and 10 mg/kg E_4 . As reference compound the anti-estro-




Figure 5: Mean (\pm SD) ultimate strength (N/mm²) at the distal femora after 4 weeks of once-daily treatment with estetrol (E_4 ; 0.1, 0.5 or 2.5 mg/(kg day)) or ethinylestradiol (EE; 0.1 mg/(kg day)) compared to vehicle in ovariectomised (OVX) rats and shamoperated controls.




Figure 6: The median percentage change in plasma levels of osteocalcin and C-telopeptide after administration of 2, 10, 20 and 40 mg/day estetrol (E_4) and 2 mg/day estradiolvalerate (E_2) for 28 days.

gen tamoxifen (TAM) was used in all 3 studies. Ovariectomy and EE at doses pharmacologically equipotent to E_4 acted as control treatments in one prevention study and in the intervention study.

A dose-dependent reduction in the number and size of tumors was seen when DMBA induced rats were co-treated with E_4 for 8 weeks, and this effect appeared to be equally effective as TAM treatment or OVX and was not seen with EE. Administration of E_4 to rats in which tumors had already been developed resulted in a significant decrease in the number and size of tumors after 4 weeks. This decrease was dose-dependent, comparable to the animals treated with TAM, and at high dose levels E_4 was as effective as OVX (Fig. 7) [24]. It was concluded that the growth of chemically induced mammary tumors is dose-dependently reduced by E_4 in female rats

and in addition has the potential to reduce also dose-dependently the number and size of already present mammary tumors. Since in these DMBA studies no blood levels of hormones (E_2 , E_4 , and gonadotrophins) were measured, the effect of E_4 observed may be explained by two different mechanisms. Either there is an antagonistic effect of E_4 in the presence of E_2 , or E_4 suppresses gonadotrophins and thereby E_2 . In the first case the effect of E_4 may be explained by antagonism at receptor level. In the second case the effect would result from suppression of E_2 , comparable to ovariectomy. Whatever the case may be, the data show that a high dose of E_4 does not stimulate tumor growth. In case E_4 would suppress E_2 the control EE would possibly do the same. However, no reduction in the number and size of tumors was seen in EE-treated animals in contrast with E_4 .




Figure 7: Intervention study. Mammary tumor count per animal (+S.E.M.) at baseline and 4 weeks after ovariectomy (OVX) or daily oral treatment with vehicle, Tamoxifen (TAM) or estetrol (E_4).

Based on this estrogen antagonistic effect, E_4 seems to be a suitable candidate for the treatment of HRT in women with (a history of) breast cancer, either for spontaneous climacteric symptoms or for symptoms induced by treatment of breast cancer with aromatase inhibitors or estrogen antagonists such as TAM. In addition E_4 may also be effective for the treatment of breast cancer itself. Currently a phase II neo-adjuvant study is ongoing investigating the effects of E_4 on proliferative and apoptotic markers in patients with breast cancer.

■ Conclusions

Estetrol is a steroid synthesized exclusively by the human fetal liver during pregnancy. After its discovery in 1965 basic research on E_4 was performed until about 1984. At that time E_4 was considered to be a weak estrogen and interest in this steroid disappeared. However, recently it has been shown that E_4 has a high and dose-related oral bioavailability in the rat [16] and the human [17], does not bind to SHBG [19] and has a long elimination half-life in both rat [16] and human [17], allowing its use as an oral once-a-day drug.

In well validated and predictive rat models E_4 behaves as an estrogen agonist in

all tissues investigated, i. e. bone [16], vagina [21], myometrium [21], endometrium [21] and brain (hot flush [20] and ovulation inhibition [20, 21]), except for breast tumor tissue where this steroid acts as an estrogen antagonist in the presence of E_2 [24].

Estetrol may be useful for a series of potential clinical applications including hormone replacement therapy in women, especially for the treatment of vaginal atrophy and hot flushes, as estrogenic component in oral contraceptives, for the prevention and treatment of osteoporosis, and E_4 might even be suitable for prevention or treatment of breast cancer. These potential applications will be further explored in clinical trials.

■ Conflict of Interest

MV is shareholder and employee of Pantarhei Bioscience, the company developing estetrol; HJT/CB is CEO and shareholder of Pantarhei Bioscience.

References:

- Hagen AA, Barr M, Diczfalussy E. Metabolism of 17 β -oestradiol-4-14C in early infancy. *Acta Endocrinol.* 1965; 49: 207–20.
- Zucconi G, Lisboa BP, Simonitsch E, Roth L, Hagen AA, Diczfalussy E. Isolation of 15 α -hydroxyl-oestriol from pregnancy urine and from the urine of newborn infants. *Acta Endocrinol* 1967; 56: 413–23.
- Schwerts J, Eriksson G, Wijkvist N, Diczfalussy E. 15 α -Hydroxylation: a new pathway of estrogen metabolism in the

human fetus and newborn, *Biochim. Biophys Acta* 1965; 100: 313–6.

4. Schwerts J, Govaerts-Videtsky M, Wijkvist N, Diczfalussy E. Metabolism of oestrone sulphate by the previable human foetus. *Acta Endocrinol* 1965; 50: 597–610.

5. Mancuso S, Benagiano G, Dell'Acqua S, Shapiro M, Wijkvist N, Diczfalussy E. Studies on the metabolism of C-19 steroids in the human foeto-placental unit. *Acta Endocrinol* 1968; 57: 208–27.

6. Heikkilä J, Adlercreutz H. A method for the determination of urinary 15 α -hydroxyestriol and estriol. *J Steroid Biochem* 1970; 1: 243–53.

7. Heikkilä J. Excretion of 15 α -hydroxyestriol and estriol in maternal urine during normal pregnancy. *J Steroid Biochem* 1971; 2: 83–93.

8. Holinka CF, Diczfalussy E, Coelingh Bennink HJT. Estetrol: a unique steroid in human pregnancy. *J Steroid Biochem Mol Biol* 2008; 110: 138–43.

9. Coelingh Bennink F, Holinka CF, Visser M, Coelingh Bennink HJT. Maternal and fetal estetrol levels during pregnancy. *Climacteric* 2008; 11 (Suppl 1): 69–72.

10. Heikkilä J, Luukkainen T. Urinary excretion of estriol and 15 α -hydroxyestriol in complicated pregnancies, *Am J Obstet Gynecol* 1971; 110: 509–21.

11. Tulchinsky D, Frigoletto FD, Ryan KJ, Fishman J. Plasma estetrol as an index of fetal well-being. *J Clin Endocrinol Metab* 1975; 40: 560–7.

12. Notation AD, Tagatz GE. Unconjugated estriol and 15 α -hydroxyestriol in complicated pregnancies. *Am J Obstet Gynecol* 1977; 128: 747–56.

13. Kundu N, Grant M. Radioimmunoassay of 15 α -hydroxyestriol (estetrol) in pregnancy serum. *Steroids* 1976; 27: 785–96.

14. Kundu N, Wachs M, Iverson GB, Petersen LP. Comparison of serum unconjugated estriol and estetrol in normal and complicated pregnancies. *Obstet Gynecol* 1981; 58: 276–81.

15. Coelingh Bennink HJT, Holinka CF, Diczfalussy E. Estetrol review: profile and potential clinical applications. *Climacteric* 2008; 11 (Suppl 1): 47–58.

16. Coelingh Bennink HJT, Heegaard AM, Visser M, Holinka CF, Christiansen C. Oral bioavailability and bone-sparing effects of estetrol in an osteoporosis model. *Climacteric* 2008; 11 (Suppl 1): 2–14.

17. Visser M, Holinka CF, Coelingh Bennink HJT. First human exposure to exogenous oral estetrol in early postmenopausal women. *Climacteric* 2008; 11 (Suppl 1): 31–40.

18. Visser M, Foidart JM, Coelingh Bennink HJT. In vitro effects of estetrol on receptor binding, drug targets and human liver cell metabolism. *Climacteric* 2008; 11 (Suppl 1): 64–8.

19. Hammond G, Hogeveen K, Visser M, Coelingh Bennink HJT. Estetrol does not bind sex hormone binding globulin or increase its production by human HepG2 cells. *Climacteric* 2008; 11 (Suppl 1): 41–6.

20. Holinka CF, Brincaat M, Coelingh Bennink HJT. Preventive effect of oral estetrol in a menopausal hot flush model. *Climacteric* 2008; 11 (Suppl 1): 15–21.

21. Heegaard AM, Holinka CF, Kenemans P, Coelingh Bennink HJT. Estrogenic uterovaginal effects of oral estetrol in the modified Allen-Doisy test. *Climacteric* 2008; 11 (Suppl 1): 22–8.

22. Coelingh Bennink HJT, Skouby S, Bouchard P, Holinka CF. Ovulation inhibition by estetrol in an in vivo model. *Contraception* 2008; 77: 186–90.

23. Coelingh Bennink HJT, Skouby S, Bouchard P, Holinka CF. Ovulation inhibition by estetrol in an in vivo model. *Climacteric* 2008; 11 (Suppl 1): 30 (a summary from *Contraception* 2008; 77: 186–90).

24. Coelingh Bennink HJT, Singer C, Simoncini T, Genazzani A, Kubista E. Estetrol, a pregnancy-specific human steroid, prevents and suppresses mammary tumor growth in a rat model. *Climacteric* 2008; 11: 29.

Mitteilungen aus der Redaktion

Besuchen Sie unsere Rubrik

[Medizintechnik-Produkte](#)



Neues CRTD Implantat
Intica 7 HF-T QP von Biotronik



Artis pheno
Siemens Healthcare Diagnostics GmbH



Philips Azurion:
Innovative Bildgebungslösung

Aspirator 3
Labotect GmbH



InControl 1050
Labotect GmbH

e-Journal-Abo

Beziehen Sie die elektronischen Ausgaben dieser Zeitschrift hier.

Die Lieferung umfasst 4–5 Ausgaben pro Jahr zzgl. allfälliger Sonderhefte.

Unsere e-Journale stehen als PDF-Datei zur Verfügung und sind auf den meisten der marktüblichen e-Book-Readern, Tablets sowie auf iPad funktionsfähig.

[Bestellung e-Journal-Abo](#)

Haftungsausschluss

Die in unseren Webseiten publizierten Informationen richten sich **ausschließlich an geprüfte und autorisierte medizinische Berufsgruppen** und entbinden nicht von der ärztlichen Sorgfaltspflicht sowie von einer ausführlichen Patientenaufklärung über therapeutische Optionen und deren Wirkungen bzw. Nebenwirkungen. Die entsprechenden Angaben werden von den Autoren mit der größten Sorgfalt recherchiert und zusammengestellt. Die angegebenen Dosierungen sind im Einzelfall anhand der Fachinformationen zu überprüfen. Weder die Autoren, noch die tragenden Gesellschaften noch der Verlag übernehmen irgendwelche Haftungsansprüche.

Bitte beachten Sie auch diese Seiten:

[Impressum](#)

[Disclaimers & Copyright](#)

[Datenschutzerklärung](#)